

In-flight transmission of foodborne disease: how can airlines improve?

Andrea Grout MSc^{1§}, Elizabeth M. Speakman MSc^{2,3}

¹James Cook University, 1 James Cook Drive, Townsville, QLD 4814, Australia

[§]Email corresponding author: andrea.grout@my.jcu.edu.au

²Edinburgh Napier University, UK

³London School of Hygiene & Tropical Medicine

Email: e.speakman@napier.ac.uk

Key Words: Epidemiology; aircraft; contamination; food hygiene; aviation food safety; densification

Abstract

Food contamination during air travel presents unique risks to those affected. Foodborne pathogens can cause serious illness among all on board, and potentially jeopardize flight safety. These risks are likely to increase with current trends of “densification” and a predicted massive expansion of air travel. While aircraft are being equipped with ever newer designs with a focus on efficiency and comfort, regulations remained largely unmodified in terms of basic hygiene requirements. Strict guidelines for food hygiene exist for on-ground food settings and catering kitchens. There is uncertainty about hygiene standards on board commercial aircraft, and little regulatory oversight of what happens to food in-flight. In two hypothetical scenarios we indicate the potential risks associated with poor food handling practice onboard aircraft, with the ultimate aim of bringing aviation food safety in line with on-ground regulations. Changes in cabin design alongside adequate training in safe food handling have the potential to increase public health protection. We urge a review of existing in-flight hygiene protocols to better direct the development of regulation, prevention, and intervention measures for aviation food safety.

Introduction

Food handling practices on board commercial aircraft are often under-regulated and there are real barriers that hinder adherence to hygiene measures. Airlines serve hundreds of millions of meals to passengers each year [1]. With the increase in global air transport, ever more people are potentially exposed to the risk of poor food hygiene in aviation settings. Due to fierce competition between airlines, there has been a growing trend of “densification”, i.e. designing aircraft to maximise seat numbers, cutting space in aircraft toilets and galleys. There are more flights, carrying more passengers, to more remote destinations and with longer flight times than ever before.

Recorded cases of food-borne disease account for only a small fraction of actual disease events [2]. The WHO estimates that each year as many as 600 million people worldwide fall ill from contaminated food, 420 000 of whom die [3]. The application of hygiene protocols is an effective measure to prevent the spread of disease [4]. Most countries have established complex, enforceable food hygiene regulations for on-ground food settings, such as ensuring that food handlers have easy access to toilets and handwashing basins. However, these regulations do not generally apply to food handling in flight and adapting standards to aircraft cabins presents a challenge: there are operational constraints, such as limited space for sanitary facilities, and also time constraints, such as having to comply with protocols and internal rules. Despite the difference with routines and rituals in on-ground food settings, food safety is governed by the same fundamental principles of hygiene, food science and public health. These disciplines have well-established theoretical foundations and robust methodologies. However, they are under-represented in the aviation environment and industry practices and are often not underpinned by enforceable legislation or lack a solid epidemiological evidence base [5].

Although aircraft are recognised as important vehicles for outbreaks and the rapid spread of foodborne diseases [6], only few reports of foodborne illness exist that are associated with aircraft [7]. This may be due to the strict food controls in airline catering stations, but many in-flight illness events go unrecognised, and may only be investigated if they have a major public health or economic impact [3]. In most instances, identification of epidemiological links between cases is extremely challenging. Illness often occurs after passengers and crew have dispersed to different public health jurisdictions [8]. Potential in-flight contamination and resulting outbreaks are difficult to differentiate from disease cases attributable to pre-

flight exposure. Outbreak investigation is further limited by ill people not seeking health care, delayed reporting, limited testing of specimens, or lack of cooperation between airlines and health authorities regarding passenger data. Even in the event of disease tracing, investigation efforts often only go back to the catering station [9]. See Box 1 for reports of outbreaks of foodborne illness associated with commercial air transport.

Box 1

Reports of outbreaks of foodborne illness associated with commercial air transport, including suspected outbreaks of Norovirus gastroenteritis from other inflight contamination sources during 1947 – 2011.

Year	Agent	Vehicle / contamination source	Origin	No. cases	Reference
1947	<i>Salmonella typhi</i>	Sandwiches	Anchorage, USA	4	[10]
1961	<i>Staphylococcus aureus</i>	Chicken	Vancouver, Canada	13	[11]
1965	<i>Staphylococcus aureus</i>	Roast turkey	Adelaide	4	[12]
1966	<i>Salmonella, staphylococcus</i>	Roast chicken	Adelaide	3	[12]
1966	<i>Staphylococcus aureus</i>	Trifle desert	New Delhi	15	[12]
1967	<i>Escherichia coli (E. coli)</i>	Oysters	London	23	[13]
1967	<i>Salmonella enteritidis</i>	Mayonnaise	Vienna	380	[12]
1969	Multiple	Unknown	Hong Kong	21	[14]
1969	Multiple	Unknown	Hong Kong	24	[15]
1970	<i>Clostridium perfringens</i>	Turkey	Atlanta, USA	25	[16]
1971	Unknown	Shrimp and crab salad	Bangkok	23	[17]
1971	<i>Shigella sonnei</i>	Unknown	Gran Canaria	219	[18]
1971	<i>Shigella sonnei</i>	Seafood cocktail	Bermuda	78	[19]
1972	<i>Vibrio parahaemolyticus</i>	Seafood appetizer	Bangkok	15	[20]
1972	<i>Vibrio cholerae</i>	Appetizer	Bahrain	47	[21]
1973	<i>Vibrio cholerae</i>	Cold asparagus & egg salad	Bahrain	66	[22]
1973	<i>Salmonella Thompson</i>	Breakfast	Denver, USA	17 (at least)	[14]
1975	<i>Staphylococcus aureus</i>	Ham	Anchorage, USA	197	[23]

1975	<i>Salmonella oranienburg</i>	Unknown	Rome	23	[12]
1976	<i>Salmonella typhimurium</i>	Cold salads	Las Palmas, Spain	550	[24, 25]
1976	<i>Salmonella brandenburg</i>	Multiple items	Paris	232	[26, 27]
1976	<i>Staphylococcus aureus</i>	Cream cakes	Rio de Janeiro	28	[28]
1976	<i>Vibrio parahaemolyticus</i>	Unknown	Bombay	28	[29]
1978	<i>Vibrio cholerae non-O1</i>	Sandwiches	Dubai	61	[29]
1982	<i>Staphylococcus aureus</i>	Custard	Lisbon	6	[14, 25]
1983	<i>Salmonella enteritidis</i>	Swiss steak	New York, USA	12	[14, 25]
1983	<i>Shigella</i> sp.	Unknown	Acapulco	42	[14, 25]
1984	<i>Salmonella enteritidis</i>	Aspic glaze	London	866	[30]
1985	<i>Salmonella enteritidis</i>	Mousse with cream	Faro	30	[31]
1986	<i>Salmonella infantis</i>	Multiple items	Vantaa	226	[32]
1988	<i>Shigella</i> sp.	Cold food items	Minnesota, USA	240	[33]
1989	<i>Salmonella enteritidis</i>	Multiple items	Spain/Finland	71	[34]
1991	<i>Salmonella</i> sp.	Unknown	Greek Island	415	[35]
1991	<i>Staphylococcus aureus</i>	Chocolate cake	Illinois, USA	26	[36]
1991	<i>Norovirus</i>	Orange juice	Melbourne	3053	[37]
1992	<i>Vibrio cholerae</i>	Seafood salad	Lima	80	[38]
1992	<i>Vibrio cholerae</i>	Seafood salad	Buenos Aires	75	[38]
1993	<i>Enterotoxigenic E. coli</i>	Unknown	Charlotte, USA	56	[39]
1997	<i>Salmonella enteritidis</i>	Chocolate éclair	Canary island	455	[40]
2002	<i>Norovirus</i> *	Contaminated surface (vomit)	London	5	[41]
2008	<i>Norovirus</i> *	Contaminated surface (vomit, faeces)	Boston, USA	22	[42]
2009	<i>Norovirus</i> *	Contaminated surface (vomit)	Unknown	27	[43, 44]
2009	<i>Norovirus</i> *	Contaminated surface (vomit)	Los Angeles, USA	63	[45]
2009	<i>Shigella sonnei</i>	Raw carrot	Hawaii	47 (at least)	[46]
2011	<i>Salmonella heidelberg</i>	Milk or eggs	Tanzania	25	[47]

* These cases were not traced to a specific food source but were likely related to other sources of contamination from inflight vomiting events. Contaminated surfaces or food preparation areas are a key transmission source for norovirus, particularly in confined spaces [48].

International air travel harbours a range of food safety hazards that emerge from the nature of aircraft cabin environments. Features of the aircraft cabin that predispose to pathogen transmission are large numbers of individuals in a confined space, and shared sanitary facilities [49]. Although the risk of in-flight food poisoning also depends on the types of foods delivered, the characteristics of people consuming the food, and the source of airline catering, contamination usually arises from unhygienic practices in food handling, inadequate food storage, and poorly enforced standards [14]. Evidence suggests that pathogens can survive for hours to months on various surfaces and spread to other individuals via direct or indirect contact. This persistence has been identified in aircraft cabins on tray tables, worktops, sink faucets and washroom door handles [50]. Larger aircraft built for longer distance and increased passenger capacity will present even greater challenges to food hygiene.

An incidence of food poisoning among crew can directly affect flight safety. For example, pilot incapacitation can have a direct impact on flight performance, and a common cause of pilot incapacitation is gastrointestinal illness [51]. Even subtle incapacitation of a pilot at a critical phase of the flight may jeopardize flight safety, such as symptoms occurring in the onset-stage of food poisoning. Regulatory and monitoring systems appear to be non-existent for in-flight food safety [52]. Few clear standards exist for hygiene requirements in aircraft cabins, and airlines generally establish their own set of cleaning standards [53]. While poor hand hygiene is often at the root of major food poisoning outbreaks, there are no requirements for a minimum number of washrooms, such as a toilet/passenger ratio, similar to an emergency door/flight attendant/passenger ratio [53], and no requirements for designated crew toilets or handwashing sinks in galleys. There is also little oversight of in-flight food handling processes, such as audits or compliance controls [52]. While aircraft are

being equipped with ever newer designs with a focus on efficiency and comfort, regulations remained largely unmodified in terms of basic hygiene requirements.

In this Commentary, we discuss three dimensions of food hygiene in-flight: onboard contamination sources, personal hygiene, and barriers to safe food handling. Two hypothetical infection scenarios illustrate the potential for in-flight contamination, aimed to highlight the divide between on-ground and in-flight food safety regulation.

Contamination Sources

Evidence suggests that about one in every five cases of food-borne illness is caused by contaminated food handlers' hands [54]. When applied to the confines of aircraft cabins, not only may contaminated hands play a key role in the occurrence of foodborne illness, but the nature of the galley design also impacts on safe food-handling practices [55]. Outbreaks of gastrointestinal illness on aircraft have been traced to in-flight incidents of vomiting in the cabin and lavatories [45]. Washroom use played a role in infection transmission when 41 travellers contracted gastrointestinal illness from one traveller's vomit [4]. The lack of recognition of vomiting events by cabin crew can lead to failure in informing destination health authorities, thereby impeding disease tracing and follow-up efforts. As passengers and crew share toilet facilities, there is a greater risk for increasing the spread of infection.

The potential for disease transmission by cabin crew is illustrated through their work in the cabin, where transmission can recur from the same source over multiple flight sectors [43]. Outbreaks resulting from indirect transmission through exposure to contaminated surfaces occurring days after the contamination incident have been reported in other contexts [56]. The type and sequence of work activity also determines the risk of contamination. For

example, failing to wash hands after touching soiled workplace surfaces is likely to be riskier than failing to wash hands after touching one's uniform. Failure to wash hands after using the toilet is likely to be riskier if the next activity is preparing a bread basket than refurbishing toiletries.

Although food handlers are typically discouraged from handling food or beverages if they have symptoms of illness that could be contagious, cabin crew were found to often fly when feeling unwell or sick [57]. Infected crewmembers may thus also act as a reservoir for disease transmission in-flight [41, 58].

Personal Hygiene and Barriers to Safe Food Handling

According to the WHO, handwashing with soap and water is the most important hygiene measure to prevent the spread of infection. There may be debate about handwashing in terms of detergents used and length of the washing process, but the benefits of handwashing in preventing foodborne illness are well documented [59]. The WHO, the International Flight Services Agency (IFSA), and the International Air Transport Association (IATA) all provide guidance on best practices on in-flight food safety and hygiene practices [1, 60, 61]. IFSA's guidance is based on the HACCP (Hazard Analysis and Critical Control Point) system, which is widely used in the food industry and which involves identification of specific hazards and measures for their control. Although the IATA notes that cabin crew should follow the same code of practice as on-ground food handlers [60], there are real barriers for crewmembers to adhere to the same stringent hand hygiene practices required for most on-ground food settings. For cabin crew to be able to apply good handwashing practice in-flight depends on (1) the number of facilities available, (2) whether handwashing facilities are in close

proximity to work stations [62] and (3) whether washrooms are vacant or galley sinks are suitable for handwashing.

Food preparation often correlates with high use of toilets by passengers (e.g. just after take-off), providing limited opportunity for crewmembers to wash their hands prior to beverage and meal service. Moreover, the combination of time pressure and lack of adequate facilities is a barrier for compliance with handwashing [63]. Cabin crew may get caught in role conflicts between safety and service tasks, which can lead to unsafe behaviour due to time constraints [64]. Similar to the way that constricted space for food handlers in small restaurants impedes adherence to good hygiene practice [65], the constraints of the aircraft galley, too, increase the risk of food safety lapses. In addition, most sinks in aircraft galleys are not designed for handwashing, as the faucet design requires one hand to operate the faucet handle [33].

There is much debate about the use of hand sanitizer products in food handling settings, with arguments such as: handwashing with soap and water is more effective for pathogen removal from hands [66, 67]; hand sanitizers should ideally be used after handwashing, but not as a substitute [68]; and hand sanitizers have no impact on hand hygiene compliance [69]. In particular, hand sanitizers are ineffective on viruses such as norovirus. Vinyl gloves can provide some protection from contamination, but they can also create a false sense of security and encourage high-risk behaviours when people are not adequately trained. Improper glove use was reported by Gaynor et al. [46] where flight catering employees touched door handles and carts with gloved hands before handling raw vegetables with gloved hands. Moreover, whether gloves can be used during service is dependent on airline-specific policy [70].

Scenarios

The following hypothetical scenarios illustrate the implications of in-flight food safety lapses, such as direct contamination by food handler hands, and opportunistic pathogen transmission through secondary sources. While these circumstances are conjectural, they represent plausible real-life events in the context of confined space conditions, limited handwashing opportunities, multitasking, role conflicts, as well as shared facilities among staff and customers. Similar to in-flight airborne disease transmission described by Han et al. [71] we assume that the movement and contact activities of cabin crew, passengers, and potentially the index case can significantly increase their personal infection risks, as well as the risk for disease transmission.

Scenario 1: Norovirus

Noroviruses are highly infectious and easily transmitted by multiple routes in confined settings, resistant to most disinfectants, and thus hard to contain using conventional sanitary measures [43]. Although typically self-limiting, severe disease cases occur in young children, the elderly, and the immunocompromised. Outbreaks of norovirus have been traced to in-flight incidents of vomiting in the cabin and lavatories [45]. On a full flight carrying 467 passengers, and a scheduled flight time of 13h 40m, a crewmember prepared four sandwich trays for premium class when she was intermittently called to the cabin for rubbish collection. Unable to wash her hands as all lavatories were occupied, she turned back to service preparations. The sandwiches were later displayed in the aircraft kitchen for self-service. Two vomiting events outside of a washroom were reported during the flight, but no disinfection of specific areas occurred. Eighteen business class passengers were part of a soccer team who resided in the same hotel as the crew during the three-day layover at the destination. Two days after arrival, vomiting and diarrhoea occurred among two crewmembers and seven

soccer players. Norovirus was confirmed as causative agent in all cases. In-flight food items were no longer available for disease tracing. Laboratory testing of retained meals at the catering kitchen showed no signs of contamination.

This scenario demonstrates the ease with which viruses can transfer between a contamination source and food items, and the potential to spread infections among people. Dissemination of norovirus is facilitated by substandard sanitary conditions and vomiting events [42], with lavatory use being a significant risk factor [59]. The pattern of norovirus outbreaks highlights the potential of aerosol transmission as well as surface contamination in confined settings [72]. Ho et al. [73] note how during a cruise ship outbreak a link could not be established to food consumption. However, the risk of gastroenteritis among passengers using shared toilet facilities was twice that of passengers who had a private facility. Consequently, the number of passengers sharing toilets was related to the rate of illness. Because 18 passengers and the crew stayed at the same location post-flight, investigative efforts were able to determine the causative agent, and to establish a likely linkage to a common contamination source. This is not usually the case. Passengers typically disperse in different directions before falling ill. Data on suspected norovirus transmission in-flight support the view that contaminated areas are rarely successfully identified and adequately treated [42, 59]. Contamination from initial vomiting events can cause infections for several days, even after routine cleaning [43, 56]. Post-flight measures dictate notification to ground staff of areas contaminated with vomit [74]. This was omitted in the scenario, implying a lack of recognition of the severity of vomiting events among crew. Only few reports of norovirus-related transmission risk exist that are associated with aircraft [45, 59, 72].

Scenario 2: Salmonella

Salmonella are resilient bacteria that can survive several weeks in dry environments and several months in water. The illness salmonellosis causes acute onset of abdominal pain, diarrhea, fever, and nausea. Children and the immunocompromised are more likely to develop severe disease. Burslem et al. [30] reported salmonella outbreaks that affected nearly 1000 passengers, aircrew and ground staff. A full flight with 352 passengers departed late. Scheduled flight time was 14h 20m. Crewmembers prepared bread baskets for premium class and stored eight hot pork dishes in the oven for sleeping passengers. Two crewmembers had been suffering from diarrhea following a previous trip but reported for work despite feeling unwell. Approximately 10 hours after the first meal was served, 12 premium class passengers, six economy class passengers, and one pilot developed symptoms of abdominal cramps and diarrhea. Five passengers and the pilot were admitted to hospital after landing. Salmonella enterotoxin was detected in all stool samples.

The source of contamination in this scenario could have been contaminated hands handling bread rolls, or inadequate storage of heated meals where bacteria multiply. In an assessment of the hygienic quality of airline meals, the most prominent contributing factors for salmonella outbreaks were found to be infected food handlers and inadequate refrigeration [75]. Salmonella bacteria have been repeatedly found in meat products [14, 76]. While bread is seen as an unusual outbreak vehicle for salmonella [77], poor personal hygiene could have contributed to the contamination. Temperatures achieved during the baking process would typically destroy any pathogen in bread, but in this scenario the bread rolls were handled after heating the bread. Delays extend the time lag between food production and consumption and increase opportunities for pathogen growth. While poor practices can involve inadequate storage at inappropriate temperatures, cabin crew may also be asymptomatic carriers of food

poisoning pathogens [78]. Travel to worldwide locations over the course of just one month puts crewmembers at heightened risk of eating or drinking contaminated food or water [52].

Discussion

Illness may not develop for days or weeks after exposure to contaminants, rendering outbreak investigation in aircraft settings extremely difficult. Passengers and crews disperse quickly, and food samples are unlikely to be available as leftover food is thrown away after a flight. Determining the real number of food poisoning incidences and contamination events on aircraft is further hampered by limited access to customer complaints and food safety-related records [52, 79].

Multi-tasking with limited access to handwashing facilities was problematic in both scenarios. Cabin crew had to smooth out service disruptions at the expense of safe handling practices. As airlines increasingly reduce space for lavatories in favour of revenue-generating seats, aircraft cabins largely remain unmodified in terms of basic hand hygiene requirements. Quantity and design of aircraft galleys and washrooms is not down to aircraft type, but to airline choice [80]. The limited space for sanitary facilities may lead to splash exposure from small wash basins, and also increase the risk of coming into contact with soiled surfaces. The scenarios underscore the importance of preventive measures such as appropriate handwashing, and proper handling and storage of food.

There is a serious lack of data regarding crew hand hygiene, or of the merits of using gloves or hand sanitisers. This presents a significant barrier to identifying the true incidence of inflight food contamination and the urgent need to evaluate the usage of provided measures such as hand sanitizers, and to adequately train crewmembers in safe food handling. While

improved hygiene may not be sufficient to break the chain of person-to-person transmission, enhanced hygiene measures are likely to reduce the transmission of norovirus during an outbreak [81]. Commercial pressures to maximise passenger numbers should not be at the expense of allowing space for adequate hygiene measures. Profits must not undermine safety. The incorporation of the internationally recognised HACCP system should become standard. Trials in the airline catering industry have been found to be cost-effective [82] and it could prove highly beneficial for onboard food safety.

Conclusion and recommendations

Food handling processes are governed by the same universal rules, whether they take place in on-ground settings or onboard aircraft. Yet attempts to contain the spread of foodborne disease via aircraft are constrained by a lack of basic hygiene infrastructure and concepts of profit over health and safety. Trends of densification mean fewer and more compact washrooms and galleys, alongside increasing passenger loads. The operation of ultra long-haul flights means increased handling of food over an extended period of time, bringing more opportunity for food safety lapses. Extended flight times also increase the risk of disease transmission and pilot incapacitation, because there is an increased risk for the sudden collapse of a crewmember resulting from food poisoning with a short incubation period.

Ensuring better adherence to in-flight food hygiene rules requires assessment of the cabin layout. Mirroring the stringent hygiene standards of on-ground food settings, there needs to be identification of those elements of the cabin layout which pose a risk to food safety and hinder personal hygiene measures. Researchers could help develop new sanitary techniques by studying what factors most influence handwashing onboard, and also look at the effectiveness of hand sanitizer gels in the cabin workspace, as well as the acceptance of hand

sanitizers by cabin crew as a substitute for handwashing. Better insight can then identify areas of weakness to design operationally feasible approaches. Airline training on hand hygiene should focus on understanding when hand hygiene is most critical, and which sanitary options are most beneficial and conducive to compliance. Developing aircraft-specific food safety plans could further serve as guidance for crew, and also raise awareness of their role as food handlers, and their importance in outbreak investigations.

Achieving onboard food safety will require a multi-pronged approach involving increased research, improved cabin design, improvements in aircrew training and behaviours, and harmonised governance. [See Box 2] The latter would require collaborative efforts of bodies such as the ICAO, IATA, IFSA and WHO. Future efforts should focus on quantifying the relative importance of in-flight disease transmission to public health. But most importantly, aircraft design should be bound to regulations that determine health and safety priorities. Just as ergonomics in galley design play an important role in preventing fatigue and injury, design should also ensure adequate handwashing opportunities. Such seemingly basic initiatives can provide a powerful means to improved food safety in aviation. Only by fixing the system of adequate facilities, regulations and inspections, and by performing the rituals of hygiene practices, can the airline industry gain the status of a 'safe' food handler.

Box 2

SUGGESTIONS TO IMPROVE ON-BOARD FOOD SAFETY

Research

- More data are required on disease transmission, including modelling and full disease tracing
- Hazard Analysis and Critical Control Point (HACCP) Analysis

Design

- Adequate sanitary facilities, e.g. sufficient toilets and wash basins

- Adequate space for good hygiene practice
- Ergonomic design, e.g. taps

Behaviours/training

- Handwashing, including use of hand sanitisers
- Food handling practice
- Management of conflicting requirements of food preparation and service provision

Governance

Collaboration between regulatory bodies to develop harmonised governance, e.g:

- Aircraft food safety plans
- Harmonised cleaning standards and policies
- Regulatory and monitoring systems

References

- [1] International Flight Services Association. World Food Safety Guidelines For Airline Catering (Version 4), http://www.ifsanet.com/?page=Food_Safety; 2016 [accessed 10 February 2019].
- [2] Scallan E, Hoeskstra RM, Angulo FJ, Tauxe RV, Widdowson, M-A, Roy SL, et al. Foodborne Illness Acquired in the United States—Major Pathogens. *Emerg Infect Dis* 2011;17(1):7-15. doi: <http://doi.org/10.3201/eid1701.P11101>.
- [3] Havelaar AH, Kirk MD, Torgerson PR, Gibb HJ, Hald T, Lake RJ, et al. on behalf of World Health Organization Foodborne Disease Burden Epidemiology Reference Group. World Health Organization Global Estimates and Regional Comparisons of the Burden of Foodborne Disease in 2010. *PLoS Med* 2015;12:e1001923. doi: <http://doi.org/10.1016/j.tmaid.2014.11.008>.
- [4] Huizer YL, Swaan CM, Leitmeyer KC, Timen A. Usefulness and applicability of infectious disease control measures in air travel: A review. *Travel Med Infect Dis* 2015;13:19-30. doi: <http://doi.org/10.1016/j.tmaid.2014.11.008>.

- [5] Grout A, Howard N, Coker R, Speakman EM. Guidelines, law, and governance: disconnects in the global control of airline-associated infectious diseases. *Lancet Infect Dis* 2017;17:e118-e122. doi: [http://doi.org/10.1016/S1473-3099\(16\)30476-5](http://doi.org/10.1016/S1473-3099(16)30476-5).
- [6] Mangili A, Gendreau MA. Transmission of infectious diseases during commercial air travel. *Lancet* 2005;365:989-96. doi: [http://doi.org/10.1016/S0140-6736\(05\)71089-8](http://doi.org/10.1016/S0140-6736(05)71089-8).
- [7] McMullan R, Edwards PJ, Kelly MJ, Millar BC, Rooney PJ, Moore JE. Food-poisoning and commercial air travel. *Travel Med Infect Dis* 2007;5(5):276-286. doi: <http://doi.org/10.1016/j.tmaid.2007.06.002>.
- [8] Enserink M. Gastrointestinal Virus Strikes European Cruise Ships. *Science* 2006;313:747. doi: <http://doi.org/10.1126/science.313.5788.747a>.
- [9] TÜV Süd. 2015. Safety considerations in airline catering, <https://www.tuv-sud.com/home-com/resource-centre/publications/e-ssentials-newsletter/food-health-e-ssentials/e-ssentials-2-2015/safety-considerations-in-airline-catering>; 2015 [accessed 8 September 2019].
- [10] Williams RB, Morley LA, Kohler M. Food-borne typhoid outbreak, with rapid dissemination of cases through air transportation. *Northwest medicine* 1950;49(10):686.
- [11] Centers for Disease Control. Staphylococcal food poisoning on a Trans-pacific Airline - Hawaii. *MMWR* 1961;10:8.
- [12] Munce B. Microbiological hazards of airline catering. *Food technology in Australia* 1978:470-476.
- [13] Preston F. An outbreak of gastro-enteritis in aircrew. *Aerospace medicine* 1968;39(5):519-521.
- [14] Tauxe RV, Tormey MP, Mascola L, Hargrett-Bean NT, Blake PA. Salmonellosis outbreak on transatlantic flights; foodborne illness on aircraft: 1947–1984. *Am J*

Epidemiol 1987;125(1):150-

7. doi: <http://doi.org/10.1093/oxfordjournals.aje.a114498.#>.

[15] Centers for Disease Control. Acute gastroenteritis among tour groups to the Orient - United States. MMWR 1969;18:301-302.

[16] World Health Organization. Gastroenteritis aboard planes. Weekly Epidemiol Rec 1971;46(15):149.

[17] Mossel D, Hoogendoorn J. Prevention of food-borne diseases in civil aviation. Indian Med Surg 1971;40(4):25-26.

[18] Oden-Johanson B, Bottiger M. Erfarenheter Fran Utbrotten av Shigella Sonnei Dysenteri Hosten 1971. (The outbreaks of Shigella sonnei dysentery in the fall of 1971). Lakartidningen 1972;69:3815-3817.

[19] Centers for Disease Control. Shigellosis related to an airplane meal - Northeastern United States. MMWR 1971;20:397-402.

[20] Peffers ASR, Bailey J, Barrow G I, Hobbs B C. Vibrio parahaemolyticus gastroenteritis and international air travel. Lancet 1973;1:143-145. doi: [http://doi.org/10.1016/s0140-6736\(73\)90207-9](http://doi.org/10.1016/s0140-6736(73)90207-9).

[21] Sutton R. An outbreak of cholera in Australia due to food served in flight on an international aircraft. Epidemiol Infect 1974;72(3):441-451. doi: <http://dodi.org/10.1017/s0022172400023688>.

[22] Dakin WP, Howell DJ, Sutton RG, O'Keefe MF, Thomas P. Gastroenteritis due to non-agglutinable (non-cholera) vibrios. Med J Austral 1974;2(13):487-490.

[23] Eisenberg MS, Gaarslev K, Brown W, Horwitz M, Hill D. Staphylococcal food poisoning aboard a commercial aircraft. Lancet 1975;306(7935):595-599. doi: [http://doi.org/10.1016/s0140-6736\(75\)90183-x](http://doi.org/10.1016/s0140-6736(75)90183-x).

- [24] World Health Organization. Salmonella typhimurium foodborne outbreak. Weekly Epidemiol Rec 1976;51:117-118.
- [25] Svensson C. Matförgiftningar på flygplan. (Food poisoning on passenger aircrafts). Svensk Veterinaertidning 1998;50:745-752.
- [26] World Health Organization. Foodborne Salmonella infections contracted on aircraft. Weekly Epidemiol Rec 1976;51(33):265-266.
- [27] Böttiger M, Romanus V. Salmonella outbreak among air passengers from Paris. Läkartidningen 1977;74(27-28):2507-2508.
- [28] World Health Organization. Outbreak of staphylococcal food poisoning aboard an aircraft. Weekly Epidemiol Rec 1976;51(51):390.
- [29] Desmarchelier P. Vibrio outbreaks from airline food and water. Food Technol Austral 1978;2:477-481.
- [30] Burslem CD, Kelly MJ, Preston FS. Food Poisoning - A Major Threat to Airline Operations. Occup Med 1990;40(3):97-100. doi: <http://doi.org/10.1093/occmed/40.3.97>.
- [31] World Health Organization. Food Safety: Microbiological quality of airline meals. Weekly Epidemiol Rec 1989;64(42):324-327.
- [32] Hatakka M. Salmonella outbreak among railway and airline passengers. Acta Vet Scand 1992;33(4):253-260.
- [33] Hedberg CW, Levine WC, White KE, Carlson RH, Winsor DK, Cameron DN, et al. An international foodborne outbreak of shigellosis associated with a commercial airline. JAMA 1992;268(22):3208-3212. doi: <http://doi.org.10.1001/jama.1992.03490220052027>.

- [34] Jahkola M. Salmonella enteritidis outbreak traced to airline food. WHO Surveillance Programme for Control of Foodborne Infections and Intoxications in Europe. Newsletter 1989;2:3.
- [35] Lambiri MA, Mavridou A, Papadakis JA. The application of hazard analysis critical control point (HACCP) in a flight catering establishment improved the bacteriological quality of meals. J Royal Soc Health 1995;115(1):26-30. doi: <http://doi.org/10.1177/146642409511500109>.
- [36] Sockett P, Ries A, Wieneke AA. Food poisoning associated with in-flight meals. Commun Dis Rep CDR Review 1993;3(7):R103-104.
- [37] Lester R, Stewart T, Carnie J, Ng S, Taylor R. Air travel-associated gastroenteritis outbreak. Commun Dis Invest 1991;15:292-93.
- [38] Eberhart-Phillips J, Besser RE, Tormey MP, Koo D, Feikin D, Araneta MR, et al. An outbreak of cholera from food served on an international aircraft. Epidemiol Infect 1996;116(1):9-13. doi: <http://doi.org/10.1017/s0950268800058891>.
- [39] Centers for Disease Control and Prevention. Foodborne outbreaks of enterotoxigenic Escherichia coli - Rhode Island and New Hampshire, 1993. MMWR 1994;43:81-89.
- [40] De Jong, B. Salmonellautbrott bland hemvavnande turister fran Kanarieoarna. Smittskydd 1998;1:6-7.
- [41] Widdowson M-A, Glass R, Monroe S, Beard RS, Bateman JW, Lurie P, et al. Probable transmission of norovirus on an airplane. JAMA 2005;293(15):1855-1860. doi:<http://doi.org/10.1001/jama.293.15.1859>.
- [42] Kirking HL, Cortes J, Burrer S, Hall AJ, Cohen NJ, Lipman H, et al. Likely transmission of norovirus on an airplane, October 2008. Clin Infect Dis 2010;50:1216-21. doi: <http://doi.org/10.1086/651597>.

- [43] Thornley CN, Emslie NA, Sprott TW, Greening GE, Rapana JP. Recurring Norovirus Transmission on an Airplane. *Clin Infect Dis* 2011;53:515-20. doi: <http://doi.org/10.1093/cid/cir465>.
- [44] Lopman B. Air sickness: vomiting and environmental transmission of norovirus on aircraft. Oxford: Oxford University Press; 2011.
- [45] Holmes J, Simmons G. Gastrointestinal illness associated with a long-haul flight. *Epidemiol Infect* 2009;137:441-7. doi: <https://doi.org/10.1017/S0950268808001027>.
- [46] Gaynor K, Park S, Kanenaka R, Colindres R, Mintz E, Ram P, et al. International foodborne outbreak of *Shigella sonnei* infection in airline passengers. *Epidemiol Infect* 2009;137:335-41. doi: <http://doi.org/10.1017/S0950268807000064>.
- [47] Rebolledo J, Garvey P, Ryan A, O'Donnell J, Cormican M, Jackson S, et al. International outbreak investigation of *Salmonella* Heidelberg associated with in-flight catering. *Epidemiol Infect* 2014;142(4):833-842. doi: <http://doi.org/10.1017/S0950268813001714>.
- [48] Brunette GW. CDC Yellow Book 2018: Health information for international travel. Oxford: Oxford University Press; 2017.
- [49] Zhao B, Dewald C, Hennig M, Bossert J, Bauer M, Pletz MW, Jandt KD. Microorganisms @ materials surfaces in aircraft: Potential risks for public health? - A systematic review. *Travel Med Infect Dis* 2019;28:6-14. doi: <http://doi.org/10.1016/j.tmaid.2018.07.011>.
- [50] Vaglenov K. Survival and Transmission of Selected Pathogens on Airplane Cabin Surfaces and Selection of Phages Specific for *Campylobacter jejuni* [dissertation]. Auburn (TX): Auburn University, <http://etd.auburn.edu/bitstream/handle/10415/4066/Corrected%20Disserta>

tion%20Vaglenov3.pdf?sequence=2&isAllowed=y; 2014 [accessed 8 September 2019].

[51] Australian Transport Safety Bureau. Analysis of Medical Conditions Affecting Pilots Involved in Accidents and Incidents from 1 January 1975 to 31 March 2006, <http://www.atsb.gov.au/media/29965/b20060170.pdf>; 2007 [accessed 8 September 2019].

[52] Sheward E. Aviation food safety. Oxford: John Wiley & Sons; 2008.

[53] International Civil Aviation Organization. Minimum Cabin Crew Requirements, <http://www.icao.int/safety/airnavigation/OPS/CabinSafety/Pages/Minimum-Cabin-Crew-Requirements.aspx>; 2017 [accessed 8 September 2019].

[54] Curtis V, Cairncross S. Effect of washing hands with soap on diarrhoea risk in the community: a systematic review. *Lancet Infect Dis* 2003;3:275-81. doi: [http://doi.org/10.1016/S1473-3099\(03\)00606-6](http://doi.org/10.1016/S1473-3099(03)00606-6).

[55] Clayton D, Griffith C, Price P, Peters A. Food handlers' beliefs and self-reported practices. *Int J Environ Health Res* 2002;12:25-39. doi: <http://doi.org/10.1080/09603120120110031>.

[56] Evans MR, Meldrum R, Lane W, Gardner D, Ribeiro C, Gallimore C, et al. An outbreak of viral gastroenteritis following environmental contamination at a concert hall. *Epidemiol Infect* 2002;129:355-60. doi: <http://doi.org/10.1017/S0950268802007446>.

[57] Mangili A, Vindenes T, Gendreau M. Infectious Risks of Air Travel. *Microbiol Spectr* 2015;3:359-66. doi: <http://doi.org/10.1128/microbiolspec.IOL5-0009-2015>.

[58] Isakbaeva ET, Widdowson MA, Beard RS, Bulens SN, Mullins J, Monroe SS, et al. Norovirus transmission on cruise ship. *Emerg Infect Dis*. 2005;11:154-8. doi: <http://doi.org/10.3201/eid1101.040434>.

- [59] World Health Organization. Guide to implementation of the WHO multimodal hand hygiene improvement strategy, <http://www.who.int/patientsafety/en/>; 2010 [accessed 10 March 2019].
- [60] International Air Transport Association. Cabin Operations Safety Best Practices Guide (Fifth Edition), https://www.iata.org/whatwedo/safety/Documents/Cabin_Operations_Safety_Best_Practices_Guide_Contents.pdf; 2019 [accessed 8 November 2019].
- [61] World Health Organization. Guide to Hygiene and Sanitation in Aviation (Third Edition), https://apps.who.int/iris/bitstream/handle/10665/44164/9789241547772_eng.pdf;jsessionid=C80E2E1455B1298698A5CC746F9F7B90?sequence=1; 2009 [accessed 8 September 2019].
- [62] Deyneko A, Cordeiro F, Berlin L, Ben-David D, Perna S, Longtin Y. Impact of sink location on hand hygiene compliance after care of patients with *Clostridium difficile* infection: a cross-sectional study. *BMC Infect Dis* 2016;16:203. doi: <http://doi.org/10.1186/s12879-016-1535-x>.
- [63] Jones AK, Cross P, Burton M, Millman C, O'Brien SJ, Rigby D. Estimating the prevalence of food risk increasing behaviours in UK kitchens. *PLoS ONE* 2017;12:e0175816. doi: <http://doi.org/10.1371/journal.pone.0175816>.
- [64] Damos DL, Boyett KS, Gibbs P. Safety versus passenger service: the flight attendants' dilemma. *Int J Aviat Psychol* 2013;23:91-112. doi: <http://doi.org/10.1080/10508414.2013.772822>.
- [65] Djekic I, Smigic N, Kalogianni EP, Rocha A, Zamioudi L, Pacheco R. Food hygiene practices in different food establishments. *Food Control* 2014;39:34-40. doi: <http://doi.org/10.1016/j.foodcont.2013.10.035>.

- [66] Foddai AC, Grant IR, Dean M. Efficacy of instant hand sanitisers against foodborne pathogens compared with hand washing with soap and water in food preparation settings: a systematic review. *J Food Prot* 2016;79:1040-54. doi: <http://doi.org/10.4315/0362-028X.JFP-15-492>.
- [67] Succo T, De Laval F, Sicard S, Belleoud D, Marimoutou C, Mayet A, et al. Do alcohol-based hand rubs reduce the incidence of acute diarrhea during military deployments? A prospective randomized trial. *Travel Med Infect Dis* 2017;15:48-51. doi: <http://doi.org/10.1016/j.tmaid.2016.11.007>.
- [68] Kampf G, Ostermeyer C. Efficacy of alcohol-based gels compared with simple hand wash and hygienic hand disinfection. *J Hosp Infect* 2004;56:13-15. doi: <http://doi.org/10.1016/j.jhin.2003.12.026>.
- [69] Muto CA, Siström MG, Farr BM. Hand hygiene rates unaffected by installation of dispensers of a rapidly acting hand antiseptic. *Am J Infect Control* 2000;28:273-6. doi: <http://doi.org/10.1067/mic.2000.103242>.
- [70] Flight Safety Foundation. Procedural Compliance Prevents Illnesses Caused by Microorganisms in Food, https://flightsafety.org/ccs/ccs_sept-oct03.pdf; 2004 [accessed 23 June 2018].
- [71] Han Z, To GNS, Fu SC, Chao CY-H, Weng W, Huang Q. Effect of human movement on airborne disease transmission in an airplane cabin: study using numerical modeling and quantitative risk analysis. *BMC Infect Dis* 2014;14:434. doi: <http://doi.org/10.1186/1471-2334-14-434>.
- [72] Kornlyo K, Kim DK, Widdowson M-A, Turabelidze G, Averhoff FM. Risk of Norovirus Transmission during Air Travel. *J Travel Med* 2009; 6:349-51. doi: <http://doi.org/10.1111/j.1708-8305.2009.00344.x>.

- [73] Ho MS, Monroe S, Stine S, Cubitt D, Glass R, Madore HP, et al. Viral gastroenteritis aboard a cruise ship. *Lancet* 1989;334:961-5. doi: [http://doi.org/10.1016/S0140-6736\(89\)90964-1](http://doi.org/10.1016/S0140-6736(89)90964-1).
- [74] Centers for Disease Control and Prevention. Preventing Spread of Disease on Commercial Aircraft: Guidance for Cabin Crew, <https://www.cdc.gov/quarantine/air/managing-sick-travelers/commercial-aircraft/infection-control-cabin-crew.html>; 2017 [accessed 8 September 2019].
- [75] Hatakka M. Hygienic quality of foods served on aircraft [dissertation]. Helsinki (Finland): University of Helsinki; 2000.
- [76] Hatakka M, Asplund K. The occurrence of Salmonella in airline meals. *Acta Vet Scand* 1993;34:391-396.
- [77] Kimura AC, Johnson K, Palumbo MS, Hopkins J, Boase JC, Reporter R, et al. Multistate Shigellosis Outbreak and Commercially Prepared Food, United States. *Emerg Infect Dis* 2004;10:1147-9. doi: <http://doi.org/10.3201/eid1006.030599>.
- [78] Cruickshank JG. Food handlers and food poisoning. *BMJ* 1990;300:207-8. doi: <https://doi.org/10.1136/bmj.300.6719.207>.
- [79] Abdelhakim A. Cabin Crew Food Safety Training: An Exploratory Study [dissertation]. Cardiff (UK): Cardiff Metropolitan University, <https://repository.cardiffmet.ac.uk/bitstream/handle/10369/8102/Ayman%20Abdelhakim%2010000757%20Final%20Draft%206%20October%202016.pdf?sequence=1&isAllowed=y>; 2016 [accessed 10 January 2019].
- [80] Airbus. Space-Flex, <https://services.airbus.com/upgrade/cabin/layout-optimisation/space-flex>; 2018 [accessed 13 November 2018].

[81] Heijne JC, Teunis P, Morroy G, Wijkmans C, Oostveen S, Duizer E, et al. Enhanced hygiene measures and norovirus transmission during an outbreak. *Emerg Infect Dis* 2009;15:24. doi: <http://doi.org/10.3201/1501.080299>.

[82] Bata D, Drosinos EH, Athanasopoulos P, Spathis P. 2006. Cost of GHP improvement and HACCP adoption of an airline catering company. *Food control* 2006;17(5):414-419. doi: <https://doi.org/10.1016/j.foodcont.2005.02.001>.