

COMMENTARY

The Ukraine crisis: Mental health resources for clinicians and researchers

Mark Shevlin¹  | Philip Hyland²  | Thanos Karatzias³ | Nino Makhshvili^{4,5} | Jana Javakhishvili⁴ | Bayard Roberts⁶

¹School of Psychology, Ulster University, Coleraine, Northern Ireland

²Department of Psychology, Maynooth University, Maynooth, Ireland

³School of Health and Social Care, Edinburgh Napier University, Edinburgh, United Kingdom

⁴Global Initiative on Psychiatry, Tbilisi, Georgia

⁵School of Business, Technology, and Education, Ilia State University, Tbilisi, Georgia

⁶Department of Health Services Research and Policy, London School of Hygiene and Tropical Medicine, London, United Kingdom

Correspondence

Mark Shevlin, Ulster University, School of Psychology, Cromore Road, Coleraine, Londonderry, BT52 1SA, Northern Ireland.

Email: m.shevlin@ulster.ac.uk

Philip Hyland, Department of Psychology, Maynooth University, Kildare, Ireland.

Email: philip.hyland@mu.ie

Abstract

The mental health consequences of the war in Ukraine will be enormous. Mental health professionals who are providing care for people in Ukraine, or those resettled elsewhere, may require access to standardized and validated assessment tools. We have developed a repository of mental health measures that are available in Ukrainian, Russian, and English and can be accessed at <http://www.traumameasuresglobal.com/ukraine>.

The world has watched in horror as the recent events in Ukraine have unfolded. After a long period of experiencing anticipatory threat as Russian troops amassed on their border, the Ukrainian population is now subjected to a full military invasion, with death, destruction, and displacement a daily reality. It almost goes without saying that the mental health consequences for the Ukrainian people will be enormous. In 2016, there were already 1,800,000 internally displaced persons (IDPs) in Ukraine (United Nations High Commissioner for Refugees, 2016), and the

Internally Displaced Persons Mental Health Survey in Ukraine (Roberts et al., 2017), carried out from March to May 2016, demonstrated that mental health problems were common among Ukrainian IDPs, with 22% of respondents reporting symptoms of depression, 18% reporting symptoms of anxiety (Roberts et al., 2019), and 55% endorsing significant levels of somatization (Cheung et al., 2018). Among current drinkers, 14.3% of men and 1.7% of women reported potentially hazardous drinking (Ramachandran et al., 2019). Moreover, the prevalence of PTSD, as assessed

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Journal of Traumatic Stress* published by Wiley Periodicals LLC on behalf of International Society for Traumatic Stress Studies.

using the *International Classification of Diseases* (11th rev.; *ICD-II*; World Health Organization [WHO], 2019) criteria, was 21% in the total sample and 57.6% among individuals experiencing clinically significant levels of impairment (Shevlin et al., 2018). To place all this in a global context, a recent study estimated the cross-national lifetime prevalence of PTSD to be 3.9% (Koenen et al., 2017). These mental health needs among IDPs in Ukraine occurred in the context of very limited mental health service availability, with 74% of participants who likely required mental health and psychosocial support care not receiving these services (Roberts et al., 2019).

At present, the Ukrainian community of mental health professionals is providing help and support in unimaginably difficult conditions; several hotlines operate, volunteers are mobilized and trained in psychological first aid provision, and mobile crisis intervention stations are in operation. However, in chaotic situations, there is a risk of offering non-evidence-based interventions, and the collection of evidence is crucially important and needs to be considered as an inevitable part of ongoing and future interventions. Resources are required for this.

The International Trauma Consortium (ITC) is a collaboration of researchers and clinicians working in the field of traumatic stress studies. To assist mental health clinicians and researchers, we have collated and hosted a range of established and validated mental health measures that have been translated into Ukrainian and Russian; these are now freely available to download at <https://www.traumameasuresglobal.com/>

The measures are:

1. The International Trauma Questionnaire (ITQ; Cloitre et al., 2018), a measure of *ICD-II* posttraumatic stress disorder (PTSD) and complex PTSD
2. The Patient Health Questionnaire-9 (PHQ-9; Kroenke et al., 2001), a brief self-report measure of depressive symptoms
3. The Generalized Anxiety Disorder-7 (GAD-7; Spitzer et al., 2006), a brief self-report measure of generalized anxiety
4. The Life Events Checklist for DSM-5 (LEC-5; Gray et al., 2004), a measure of trauma exposure
5. The Patient Health Questionnaire-15 (Kroenke et al., 2002), a measure of somatic symptom severity
6. The World Health Organization Disability Assessment Schedule-II (WHODAS-II; WHO, 2001), a brief measure of functional impairment

We hope that these measures are helpful to clinicians and researchers who will be dealing with the psychological aftermath of the conflict in Ukraine. We intend to continue to expand this repository—for example, child and adoles-

cent measures will soon be added—and we would welcome any other resources (email Mark Shevlin or Philip Hyland).

ORCID

Mark Shevlin  <https://orcid.org/0000-0001-6262-5223>

Philip Hyland  <https://orcid.org/0000-0002-9574-7128>

REFERENCES

- Cheung, A., Makhshvili, N., Javakhishvili, J., Karachevsky, A., Kharchenko, N., Shpiker, M., & Roberts, B. (2019). Patterns of somatic distress among internally displaced persons in Ukraine: Analysis of a cross-sectional survey. *Social Psychiatry and Psychiatric Epidemiology*, 54(10), 1265–1274. <https://doi.org/10.1007/s00127-019-01652-7>
- Cloitre, M., Shevlin, M., Brewin, C. R., Bisson, J. I., Roberts, N. P., Maercker, A., Karatzias, T., & Hyland, P. (2018). The International Trauma Questionnaire: Development of a self-report measure of *ICD-II* PTSD and complex PTSD. *Acta Psychiatrica Scandinavica*, 138(6), 536–546. <https://doi.org/10.1111/acps.12956>
- Gray, M., Litz, B., Hsu, J., & Lombardo, T. (2004). Psychometric properties of the Life Events Checklist. *Assessment*, 11(4), 330–341. <https://doi.org/10.1177/1073191104269954>
- Koenen, K., Ratanatharathorn, A., Ng, L., McLaughlin, K., Bromet, E., Stein, D., Karam, E. G., Meron Ruscio, A., Benjet, C., Scott, K., Atwoli, L., Petukhova, M., Lim, C. C. W., Aguilar-Gaxiola, S., Al Hamzawi, A., Alonso, J., Bunting, B., Ciutan, M., de Girolamo, G., Degenhardt, L., ... Kessler, R. (2017). Posttraumatic stress disorder in the World Mental Health Surveys. *Psychological Medicine*, 47(13), 2260–2274. <https://doi.org/10.1017/S0033291717000708>
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16(9), 606–613. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x>
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2002). The PHQ-15: Validity of a new measure for evaluating the severity of somatic symptoms. *Psychosomatic Medicine*, 64(2), 258–266. <https://doi.org/10.1097/00006842-200203000-00008>
- Ramachandran, A., Makhshvili, N., Javakhishvili, J., Karachevskyy, A., Kharchenko, N., Shpiker, M., Ezard, N., Fuhr, D. C., & Roberts, B. (2019). Alcohol use among conflict-affected persons in Ukraine: Risk factors, coping and access to mental health services. *European Journal of Public Health*, 29(6), 1141–1146. <https://doi.org/10.1093/eurpub/ckz117>
- Roberts, B., Makhshvili, N., & Javakhishvili, J. (2017). *Hidden burdens of conflict: Issues of mental health and access to services among internally displaced persons in Ukraine*. <https://www.international-alert.org/publications/hidden-burdens-of-conflict/>
- Roberts, B., Makhshvili, N., Javakhishvili, J., Karachevskyy, A., Kharchenko, N., Shpiker, M., & Richardson, E. (2019). Mental health care utilisation among internally displaced persons in Ukraine: Results from a nation-wide survey. *Epidemiology and Psychiatric Sciences*, 28(1), 100–111. <https://doi.org/10.1017/S2045796017000385>
- Shevlin, M., Hyland, P., Vallières, F., Bisson, J., Makhshvili, N., Javakhishvili, J., Shpiker, M., & Roberts, B. (2018). A comparison of *DSM-5* and *ICD-II* PTSD prevalence, comorbidity and disability: An analysis of the Ukrainian Internally Displaced Person's

- Mental Health Survey. *Acta Psychiatrica Scandinavica*, 137(2), 138–147. <https://doi.org/10.1111/acps.12840>
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*, 166(10), 1092–1097. <https://doi.org/10.1001/archinte.166.10.1092>
- United Nations High Commissioner for Refugees. (2016). *Global trends: Forced displacement in 2016*. <http://www.unhcr.org/globaltrends2016/>
- World Health Organization. (2019). *International statistical classification of diseases and related health problems* (11th rev.). <https://icd.who.int/>
- World Health Organization. (2001). *World Health Organization Disability Assessment Schedule II*. [https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-\(-whodas-2.0\)/](https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-(-whodas-2.0)/)

How to cite this article: Shevlin, M., Hyland, P., Karatzias, T., Makhshvili, N., Javakhishvili, J., & Roberts, B. (2022). The Ukraine crisis: Mental health resources for clinicians and researchers. *Journal of Traumatic Stress*, 1–3. <https://doi.org/10.1002/jts.22837>