

Short communication

Definitive Glasgow acute pain scale for cats: validation and intervention level

J. Reid, E. M. Scott, G. Calvo, A. M. Nolan

THE usefulness of a pain assessment instrument is enhanced in general practice if the score can be linked to an intervention level, which is informative as to whether or not an animal requires analgesic treatment. Previously the authors described the derivation of an intervention level for the Glasgow Composite Measure Pain Scale (CMPS) short form tool for assessing acute pain in dogs (Reid and others 2007). More recently, the authors reported the validation of a behaviour-based tool (the rCMPS-F) for the assessment of acute pain in cats that was developed using psychometric principles (Calvo and others 2014). It takes the form of a structured questionnaire completed by an observer following a standard protocol and includes assessment of spontaneous and evoked behaviours, interactions with the animal and clinical observations. Construction and clinical testing of the tool supported its validity and provided some evidence for responsiveness, but sensitivity was moderate (misclassification, 26.7 per cent). In order to improve the performance of the tool, a simple three-point facial scale, which in preliminary testing had performed very well in classifying cats in pain, was developed (Holden and others 2014) with the intention of embedding it within the behaviour-based cat tool, an approach adopted previously in paediatric medicine (CRIES (Krechel and others 1995); Premature Infant Pain Profile (Stevens and others 1996)). This communication describes a single, multicentre study designed to (i) validate the combined tool (CMPS-Feline) using a known groups design based on those cats requiring analgesia compared with those that did not according to clinical veterinary impression and (ii) derive an intervention-level score for analgesia for the composite tool, the CMPS-Feline.

The facial component was embedded within the rCMPS-F as shown in Fig 1. In total, 119 cats, 65 males and 54 females, median age 63 months (range 1–240 months) undergoing postoperative care or having been admitted for surgery or any acutely

painful trauma or medical condition to one of three University Veterinary Schools, four small animal general practices or two charity veterinary clinics were recruited to the study. No restrictions were placed on the breed, age or sex of the cats, the type of surgical procedure, trauma or medical condition or timing of the assessment. All cats were evaluated for sedation using a sedation scoring scale (0–3) modified from Lascelles and others (1994) and previously described (Calvo and others 2014); any cats with a score >1 were excluded from the study. In total, 45 cats underwent surgical procedures and 74 did not. All cats were scored by a veterinary nurse/technician using the CMPS-Feline and thereafter a veterinary surgeon, blinded to the score responded to the question 'Do you think this animal requires analgesia? Yes/no'. The total score (from a possible maximum of 20) was computed by summing the answers to all questions. The project was approved by the University of Glasgow Veterinary School ethics committee.

Boxplots were used to compare cats that required analgesia compared with those that did not. Formal analysis involved Mann-Whitney and 95 per cent CIs for medians. Linear discriminant analysis (LDA)¹ was used to determine the intervention-level score. Statistical analyses were carried out using MINITAB version 14 (Minitab, Microsoft Corporation). Of the 119 cats, 49 were recorded as not requiring analgesia, 70 were recorded as requiring analgesia with median scores 1 (range 0–8) and 8 (range 2–16) respectively for the two groups (Fig 2). LDA was first used to classify cats as requiring analgesia (yes/no). Using the total score from the combined tool, 82.4 per cent of the cats were correctly classified with the intervention level set at 5 and above. From the results, 43 out of 49 (88 per cent) of the no analgesia cats and 55 out of 70 (78.6 per cent) of analgesia cats were correctly identified.

Hypotheses used for testing construct validity of pain scales include 'known groups' validity where the instrument should be able to distinguish correctly between groups expected to have different scores. Facial expression is considered a sensitive indicator of noxious procedures, and extensive research has gone into its use for measuring acute and postoperative pain intensity in neonates (Grunau and others 1998, Tomlinson and others 2010). Work conducted in animals has led to the development of standardised facial coding systems for recording pain in rodents, for example, the Mouse Grimace Scale (Langford and others 2010); Rat Grimace Scale (Sotocinal and others 2011), rabbits (Keating and others 2012) and, more recently, a pain expression scale for horses was described (Dalla Costa and others 2014). Facial expression scales have been previously incorporated into multidimensional measure pain instruments (Stevens and others 1996, Hand and others 2010), and the authors hypothesised that the incorporation of their simple facial scale would improve the performance of the behaviour-based tool. The result from this study has indicated enhanced discriminatory ability of the holistic tool, the CMPS-Feline, compared with the rCMPS-F (misclassification rates of 17.6 per cent and 26.7 per cent, respectively). Brondani and others (2013) defined an intervention level for the English version of their cat pain tool; however, there were marked differences in study designs compared with this work. All their cats underwent the same standardised soft tissue procedure (ovariohysterectomy) and scoring was performed by observers trained in anaesthesia. In contrast, the study reported here was undertaken in clinical environments where acute pain would arise

Veterinary Record (2017)

doi: 10.1136/vr.104208

J. Reid,
School of Veterinary Medicine, University
of Glasgow, Glasgow G61 1QH, UK
E. M. Scott,
School of Mathematics & Statistics,
University of Glasgow, Glasgow G12
8QW, UK
G. Calvo,
Royal Veterinary College, University
of London, London AL9 7TA, UK

A. M. Nolan,
Edinburgh Napier University,
Edinburgh EH11 4BN, UK
E-mail for correspondence: a.
nolan@napier.ac.uk
Provenance: not commissioned;
externally peer reviewed
Accepted January 2, 2017

¹Discriminant analysis creates a classification rule that can be used to predict which class or group (analgesia yes/no) a cat belongs to. This is done by estimating the probabilities that the cat belongs to each class/group.

Glasgow Feline Composite Measure Pain Scale: CMPS- Feline

Choose the most appropriate expression from each section and total the scores to calculate the pain score for the cat. If more than one expression applies choose the higher score

LOOK AT THE CAT IN ITS CAGE:

Is it?

Question 1

- Silent / purring / meowing 0
- Crying / growling / groaning 1

Question 2


- Relaxed 0
- Licking lips 1
- Restless/cowering at back of cage 2
- Tense/crouched 3
- Rigid/hunched 4

Question 3

- Ignoring any wound or painful area 0
- Attention to wound 1

Question 4

- a) Look at the following caricatures. Circle the drawing which best depicts the cat's ear position?



- b) Look at the shape of the muzzle in the following caricatures. Circle the drawing which appears most like that of the cat?




FIG 1: The Glasgow Composite Measure Pain Scale-Feline

from varied sources including postsurgical, trauma and medical cases, and where its assessment would be undertaken by observers of varying levels of experience, thus assessing its robustness for use in general practice. In summary, the CMPS-Feline with its embedded facial image component has been shown to be a

valid scale for the measurement of acute pain in cats in general veterinary practice with a recommended intervention level of 5 and above (out of a total possible score of 20). The CMPS-Feline displayed increased discriminatory ability over the previously described rCMPS-F.

285 **APPROACH THE CAGE, CALL THE CAT BY NAME & STROKE ALONG ITS BACK** 356
 286 **FROM HEAD TO TAIL** 357

287 **Question 5** 358

288 Does it? 359

289 Respond to stroking 0 360

290 Is it? 361

291 Unresponsive 1 362

292 Aggressive 2 363

293 **IF IT HAS A WOUND OR PAINFUL AREA, APPLY GENTLE PRESSURE 5 CM** 364
 294 **AROUND THE SITE. IN THE ABSENCE OF ANY PAINFUL AREA APPLY SIMILAR** 365
 295 **PRESSURE AROUND THE HIND LEG ABOVE THE KNEE** 366

296 **Question 6** 367

297 Does it? 368

298 Do nothing 0 369

299 Swish tail/flatten ears 1 370

300 Cry/hiss 2 371

301 Growl 3 372

302 Bite/lash out 4 373

303 **Question 7** 374

304 General impression 375

305 Is the cat? 376

306 Happy and content 0 377

307 Disinterested/quiet 1 378

308 Anxious/fearful 2 379


309 Dull 3 380

310 Depressed/grumpy 4 381

311 **Pain Score ... /20** 382

312 © Universities of Glasgow & Edinburgh Napier 2015. Licensed to NewMetrica Ltd. Permission granted to 383
 313 reproduce for personal and educational use only. To request any other permissions please contact 384
 314 jacky.reid@newmetrica.com. 385

315 **FIG 1 Continued** 386



317 **FIG 2: Boxplot of known group CMPS-Feline pain scores.** 388
 318 **Mann-Whitney P value <0.005 and 95 per cent CI for difference in** 389
 319 **median (analgesia (yes)-analgesia (no)) was 6–8** 390

320 Twitter Follow Andrea Nolan @ProfAndreaNolan 391

321 **Acknowledgements** 392

322 The authors are grateful to all colleagues in veterinary practice, 393
 323 the Scottish SPCA, PDSA and University Veterinary Schools 394
 324 who contributed to the success of this study, especially Eleanor 395
 325 Holden and Andrew Bell (University of Glasgow) and Sheilah 396
 326 Robertson (Michigan State University). 397

327 **References** 398

- 328 BRONDANI, J. T., MAMA, K. R., LUNA, S. P., WRIGHT, B. D., NIYOM, S., 399
 329 AMBROSIO, J., VOGEL, P. R. & PADOVANI, C. R. (2013) Validation of the 400
 330 English version of the UNESP-Botucatu multidimensional composite pain scale 401
 331 for assessing postoperative pain in cats. *BMC Veterinary Research* 9, 143 402
 332 CALVO, G., HOLDEN, E., REID, J., SCOTT, E. M., FIRTH, A., BELL, A., 403
 333 ROBERTSON, S. & NOLAN, A. M. (2014) Development of a behaviour-based 404
 334 measurement tool with defined intervention level for assessing acute pain in cats. 405
 335 *Journal of Small Animal Practice* 55, 622–629 406
 336 DALLA COSTA, E., MINERO, M., LEBELT, D., STUCKE, D., CANALI, E., & 407
 337 LEACH, M. C. (2014) Development of the Horse Grimace Scale (HGS) as a pain 408
 338 assessment tool in horses undergoing routine castration. *PLOS ONE* 9, e92281 409
 339 GRUNAU, R. E., OBERLANDER, T., HOLSTI, L. & WHITFIELD, M. F. (1998) 410
 340 Bedside application of the Neonatal Facial Coding System in pain assessment of 411
 341 premature neonates. *Pain* 76, 277–286 412
 342 HAND, I. L., NOBLE, L., GEISS, D., WOZNIAK, L. & HALL, C. (2010) COVERS 413
 343 neonatal pain scale: development and validation. *International Journal of Pediatrics* 414
 344 2010, 496719 415
 345 HOLDEN, E., CALVO, G., COLLINS, M., BELL, A., REID, J., SCOTT, E. M. & 416
 346 NOLAN, A. M. (2014) Evaluation of facial expression in acute pain in cats. 417
 347 *Journal of Small Animal Practice* 55, 615–621 418
 348 KEATING, S. C. J., THOMAS, A. A., FLECKNELL, P. A. & LEACH, M. C. (2012) 419
 349 Evaluation of EMLA cream for preventing pain during tattooing of rabbits: 420
 350 changes in physiological, behavioural and facial expression responses. *PLoS ONE* 421
 351 7, e44437 422
 352 KRECHEL, S. W., & BILDNER, J. (1995) CRIES: a new neonatal postoperative pain 423
 353 measurement score. Initial testing of validity and reliability. *Paediatric Anaesthesia* 424
 354 5, 53–61 425
 355 LANGFORD, D. J., BAILEY, A. L., CHANDA, M. L., CLARKE, S. E., 426
 356 DRUMMOND, T. E., ECHOLS, S., GLICK, S., INGRAO, J., KLASSEN-ROSS, T., 427
 357 LACROIX-FRALISH, M. L., MATSUMIYA, L., SORGE, R. E., SOTOCINAL, 428
 358 S. G., TABAKA, J. M., WONG, D., VAN DEN MAAGDENBERG, A. M. J. M., 429
 359 FERRARI, M. D., CRAIG KENNETH, D. & MOGIL, J. S. (2010) Coding of facial 430
 360 expressions of pain in the laboratory mouse. *Nature Methods* 7, 447–449 431
 361 LASCELLES, B. D. X., BUTTERWORTH, S. J. & WATERMAN, A. E. (1994) 432
 362 Postoperative analgesic and sedative effects of carprofen and pethidine in dogs. 433
 363 *Veterinary Record* 134, 187–191 434
 364 REID, J., NOLAN, A. M., HUGHES, J. M. L., LASCELLES, D., PAWSON, P. & 435
 365 SCOTT, E. M. (2007) Development of the short-form Glasgow Composite 436
 366 Measure Pain Scale (CMPS-SF) and derivation of an analgesic intervention score. 437
 367 *Animal Welfare* 16, 97–104 438

427	SOTOCINAL, S. G., SORGE, R. E., ZALOUM, A., TUTTLE, A. H., MARTIN, L. J.,	TOMLINSON, D., VON BAEYER, C. L., STINSON, J. N. & SUNG, L. (2010) A	498
428	WIESKOPF, J. S., MAPPLEBECK, J. C. S., WEI, P., ZHAN, S., ZHANG, S.,	systematic review of faces scales for the self-report of pain intensity in children.	499
429	MCDUGALL, J. J., KING, O. D., & MOGIL J. S. (2011) The Rat Grimace Scale:	<i>Pediatrics</i> 126, e1168–e1198.	500
Q5430	a partially automated method for quantifying pain in the laboratory rat via facial		501
431	expressions. <i>Molecular Pain</i> 7, 55		502
432	STEVENS, B., JOHNSTON, C., PETRYSHEN, P., & TADDIO, A. (1996) Premature		503
433	Infant Pain Profile: development and initial validation. <i>The Clinical Journal of Pain</i>		504
434	12, 13–22		505
435			506
436			507
437			508
438			509
439			510
440			511
441			512
442			513
443			514
444			515
445			516
446			517
447			518
448			519
449			520
450			521
451			522
452			523
453			524
454			525
455			526
456			527
457			528
458			529
459			530
460			531
461			532
462			533
463			534
464			535
465			536
466			537
467			538
468			539
469			540
470			541
471			542
472			543
473			544
474			545
475			546
476			547
477			548
478			549
479			550
480			551
481			552
482			553
483			554
484			555
485			556
486			557
487			558
488			559
489			560
490			561
491			562
492			563
493			564
494			565
495			566
496			567
497			568

