

Note: Snapshot PDF is the proof copy of corrections marked in EditGenie, the layout would be different from typeset PDF and EditGenie editing view.

## Author Queries & Comments:

**Q1** : Please check whether the doctype has been set correctly

Response: Resolved

**Q2** : Please provide a short biography of the author(s).

Response: Yvonne Kuipers, Professor of Midwifery; Julie DeGrave, midwife researcher; Valerie Bosmans, midwife researcher; Ellen Thaelts, RM and PhD student; Eveline Mestdagh, Head of Midwifery Education

**Q3** : Please provide complete affiliation details including postal/zip code to be included in the contact field.

Response: Resolved

**Q4** : The abstract is currently too long. Please edit the abstract down to no more than 200 words.

Response: Resolved

**Q5** : Please check whether the P values represented in the sentence "Quantitative data were..." are correct.

Response: These p-values are correct.

**Q6** : Please check whether the formatting given in table 3 and its footnotes are correct.

Response: Resolved

**Q7** : The disclosure statement has been inserted. Please correct if this is inaccurate.

Response: Resolved

**Q8** : The CrossRef database (www.crossref.org/) has been used to validate the references. Mismatches between the original manuscript and CrossRef are tracked in red font. Please provide a revision if the change is incorrect. Do not comment on correct changes.

Response: Resolved

**Q9** : Please provide missing page range for reference "[16]" references list entry.

Response: No page range available

**Q10** : Please provide missing page range for reference "[21]" references list entry.

Response: Resolved

**Q11** : Please provide missing page range for reference "[29]" references list entry.

Response: Resolved

**Q12** : Please provide missing page range and editors name for reference "[42]" references list entry.

Response: Resolved

**Q13** : Please provide missing volume number for reference "[83]" references list entry.

Response: Resolved

**Q14** : Please provide missing page range for reference "[92]" references list entry.

Response: No page numbers available

**Q15** : Please provide missing page range for reference "[96]" references list entry.

Response: Resolved

REVIEW ARTICLE

# Midwifery-led care[Q1]: A single mixed-methods synthesis

**Recto running head** : INTERNATIONAL JOURNAL OF HEALTHCARE MANAGEMENT

**Verso running head** : Y. KUIPERS ET AL.



 Yvonne Kuipers<sup>a,b,c</sup>,  Julie Degraeve<sup>a,c</sup>,  Valerie Bosmans<sup>a</sup>,  Ellen Thaelts<sup>d</sup>,  Eveline Mestdagh<sup>a,c</sup> [Q2]

<sup>a</sup> Department of Health and Social Care, School of Midwifery, AP University of Applied Sciences, Antwerp, Belgium

<sup>b</sup> School of Health and Social Care, Edinburgh Napier University, Edinburgh, UK

<sup>c</sup> Faculty of Medicine & Health Sciences, Department of Nursing and Midwifery, Antwerp University, Wilrijk, Belgium

<sup>d</sup> Faculty of Health & Wellbeing, School of Community Health & Midwifery, University of Central Lancashire, Preston, UK

**CONTACT** Yvonne Kuipers,  Edinburg Napier University, Sighthill Campus, Edinburgh EH11 4BN, Scotland  y.kuipers@napier.ac.uk yvonne.fontein-kuipers@ap.be [Q3]

**History** : received : 2022-2-3 accepted : 2022-4-21

**Copyright Line** : © 2022 Informa UK Limited, trading as Taylor & Francis Group

## ABSTRACT

### Background

Although midwifery-led care (MLC) has shown to be beneficial for women and their children and for midwives, the implementation of MLC remains challenging. The midwife is the central care provider in MLC. An exploration of the potential factors that might attribute to midwives' utility of MLC will offer relevant information to support MLC implementation, scale up, and MLC sustainability. Midwifery Led Care (MLC) has shown to be beneficial for women and for midwives. The implementation of MLC remains challenging.

### Objective

To explore the utility of MLC and midwives' MLC behavioural determinants.

## Methods

A systematic mixed-methods review was conducted, integrating data derived from methodologically different studies into a single mixed-methods synthesis. Quality was assessed using The Joanna Briggs Institute Critical Appraisal tools. Data were organized using the Feasibility–Appropriateness–Meaningfulness–Effectiveness (FAME) scale. Behavioural MLC determinants were grouped in an extended Attitude–Social influence–Self-efficacy (ASE) model. After a synthesis and summary of the data and a descriptive thematic analysis, all FAME/ASE variables were quantified for a Bayesian Pearson correlation analysis of the FAME and ASE themes. The Bayes Factor (BF) indicated the strength between the correlations.

## Results

Twenty-six papers of good quality were included. The relationships between the FAME scales and ASE themes showed very strong evidence (BF 31.1–41.6), strong (BF 11.2–28.5) and to a lesser degree moderate (BF 3.1–9.7), and anecdotal evidence (BF 1.5–2.9). MLC utility was predominantly explained by the appropriateness and effectiveness of MLC and their correlations with the midwife's attitude, the perceived social influence of the public, supportive factors, regulation, professional and personal norms, and intention. Anecdotal to very strong evidence for correlations between the ASE themes (BF 1.5–41.6) was observed.

## Conclusion

To implement, scale up, and maintain MLC, a multipronged approach is needed, with attention for the strong and very strong behavioural aspects of midwives related to the utility of MLC. To fully embed MLC as a first choice for women, midwives need to stand up for their professional identity in the wider culture and climate of maternity services to push the change for MLC [Q4].

## KEYWORDS

- Behaviour
- care management
- continuity of care
- midwifery organization
- organizational culture
- review
- midwife-led care
- caseload midwifery

## FUNDING

This work was supported by Department of Economy, Science, and Innovation of the Flemish Government [grant number PWO-3105R2007].

## Introduction

Midwife-Led Care (MLC) is a care model where the midwife is the lead professional in planning, organizing, and providing care to a woman from booking to the postnatal period within a multi-disciplinary network of consultation and referral with other care providers [1]. MLC is widely recommended as a measure of quality-of-care since it has shown to improve outcomes such as the reduction of intrapartum medical interventions, lower rates of preterm birth, intrapartum pain relief, amniotomy, and episiotomy, and higher rates of spontaneous vaginal birth and breastfeeding, as well as higher rates of care satisfaction among women and job satisfaction among midwives [1–10]. The philosophy behind MLC is the normality of childbirth, continuity of care and being cared for by a known, trusted midwife during labour, aiming to optimize bio-psychosocial processes, strengthening the opportunities for women to achieve a physiological birth and a positive birth experience [1,11–13].

There is great diversity in how MLC is organized across the different healthcare settings, such as community practices, hospitals, freestanding, or alongside birth units applying various care strategies [14]. Examples of MLC strategies are continuity of care from a known midwife, caseload midwifery, continuity of care management, and team/group-midwifery [1,2,12,13,15–18]. Despite good quality evidence, MLC is still not utilized as a worldwide standard model of care [19–22]. This, despite the International Confederation of Midwives (ICM) [23] states that MLC should be the first choice for all childbearing women.

Midwives are the essential caregivers and primary executors at the heart of MLC. Given the midwifery scope and the pivotal role of midwives within MLC, an understanding of midwives' behaviour is of extreme importance to shape the development of MLC [9,11,22,24] – that is, 'what makes midwives tick' to provide MLC, what helps and hinders them, how do they (inter)act as MLC midwives, how do they respond to and view MLC. To create awareness about how to increase the (transformational) utility of MLC, an exploration of midwives' behavioural factors possibly playing a role in implementation, transition, capacity building, and evaluation processes of MLC is essential [25–29]. A systematic approach to explore midwives' behaviour can contribute to a synthesis of what is known and what needs to be elaborated on regarding the context or environment, organizational culture, (recurring) patterns, beliefs, and values of midwives about MLC and how and with whom the midwife (inter)acts [26]. So far, there is no framework that systematically focuses on the relation between the behaviour of midwives and MLC utility to identify strengths or potential weaknesses of MLC, to inform practice, and/or to evaluate the MLC model. Midwives, those who already work within an MLC model of care or who execute MLC strategies, those who are scoping or anticipating MLC practices, as well as midwifery managers and educators, would benefit from evidence on possible implementation strategies for MLC. Information is needed as to whether MLC is a model that is feasible, appropriate, adequate, and meaningful in maternity services.

The present study aims to explore the utility and behavioural determinants of performing MLC reported by midwives and to examine the relationship between the behaviour of midwives and MLC utility. It was hypothesized that midwives' behaviour is correlated with the utility of MLC. A framework of MLC was proposed, integrating and building upon previous research by integrating multilevel variables of multilevel research, with varying sample sizes, allowing the operationalization of theoretical thinking and creation of theory [30,31].

## Methods

### Design

A systematic mixed-methods review was performed, integrating data derived from methodologically different studies into a single mixed-methods synthesis [32]. A synthesis of the data, derived from the review, was conducted, data entries were summarized, the patterns and commonalities emerging from the data were described, and the relationships between the findings were analysed [33].

### Search strategy


The literature search was conducted (JD, VB) between November 2020 and May 2021 in the following electronic databases: PubMed, ScienceDirect, Web of Science, Scopus, PsychINFO, and Biomed Central. The following key terms and database-specific advanced search options were used: (midwife) AND (model of care OR continuity of care OR models, organizational) AND (midwives experience OR midwives' perception OR midwives' satisfaction OR attitude of health personnel OR barriers). A 10-year limit was placed on publication dates. Both original qualitative and quantitative literature published in the English language were included.

### Selection

The purpose was to select literature exploring either midwives' reports working in the MLC model or midwives' perceptions of the hypothetical or potential (future) utility of

the model. Primary peer-reviewed studies with samples of practising midwives in countries enlisted by the ICM [34] were included. Studies with healthcare professionals other than midwives, studies unclear if or what data were obtained from midwives, and studies showing ambiguity whether the midwife enacted as primary care giver was excluded. From this search, additional publications were identified via the references of systematic reviews and forward citations. Of the 290 records identified, 199 remained after removing the duplicates and papers that did not pertain to the focus of this review (JD, VB). Following this, 56 publications remained and were screened for their eligibility against the inclusion and exclusion criteria, leaving 30 records prior to quality appraisal (Figure 1).

**Figure 1.** PRISMA 2020 flow diagram. From: Page et al. [35].



## Quality appraisal

The 30 studies were appraised using The Joanna Briggs Institute (JBI) Critical Appraisal tools for use in systematic reviews (YK, JD, VB). A scoring system was added to this process to assist in summarizing the quality level. Each item was rated: 'Yes', 'No', 'Unclear', or 'Not/Applicable'. Based on the overall appraisal of items, the paper was included or excluded or further explored and discussed [36,37]. The components of studies with a mixed-methods design were separately appraised. The papers with two or more absent and/or unclear JBI-criteria were discussed. After reaching consensus (YK, JD, VB), four papers were excluded, leaving a final 26 good quality papers for further synthesis and analysis (Figure 1).

## Organizing the data

To organize the data and to analyse the relevant literature about MLC utility, the Feasibility–Appropriateness–Meaningfulness–Effectiveness (FAME) scale was used[33]. In this study, feasibility is about whether MLC is socio-politically and financially possible, regulated, and managed within the climate of maternity services and the work–life balance from an employee/employer viewpoint. Appropriateness is how MLC relates to the midwife's professional and personal domain, culture, and context. Meaningfulness is about how midwives make sense of MLC, the purposefulness and worth of MLC, relying on personal experiences, opinions, values, beliefs, and thoughts. Effectiveness is about the relationship between MLC and (clinical or health) outcomes, including the satisfaction of women and of midwives[33]. To organize the behavioural determinants of MLC, the Attitude–Social influence–self-efficacy (ASE) model was chosen. According to this model, a person's specific behaviour is a result of several influencing cognitive factors. The intention to perform a certain behaviour predicts the actual behaviour – in this case, MLC. The three central psychosocial determinants attitude, social influence, and self-efficacy are assumed to primarily predict and determine the midwife's intention to provide MLC. Intention can also be influenced by external variables, such as supportive factors, barriers, and/or knowledge [38]. Earlier studies have used the (extended) ASE model to explain midwifery practice and midwives' behaviour [39–41].

## Data extraction & synthesis

The aim of the synthesis was to summarize and synthesize the reports of midwives' reality as represented by the primary authors. First, all 26 articles were (re)read, focusing on the content of the data sources (YK, JD, VB). Because the research question focused on midwives' reports, the data was extracted from the studies' results sections. After immersing in the data, per paper, line by line, relevant ASE variables (quantitative studies) and ASE text segments (qualitative studies) were identified, assigning a code fitting an ASE category[42]. Data-driven additional behavioural aspects, such as professional, personal, and social norms, regulation, enablers, supportive factors, barriers, knowledge, and skills were added to the original ASE model to provide a full scope of midwives' behaviour[32,43,44]. These additional behavioural concepts were integrated to increase the predictiveness of MLC[31,32]. A matrix (Microsoft Excel®) was created to group the ASE codes and categories. Similar categories were grouped together (e.g. supportive and enabling factors, knowledge and skills), resulting in 12 final ASE themes[42]. The ASE themes were then ordered in the four FAME scales; each column of the ASE themes corresponded with the rows of the FAME scale. Findings were compared and refined by all authors until consensus was reached about the relevant place in the matrix. Second, patterns in the FAME and the ASE themes were identified across the text segments and the variables within and between the entries. This was an iterative and recursive process of constant comparison and contrasting between all authors to establish a level of focus regarding the FAME and ASE themes[45]. The viewpoints were presented as themes [42]. The number of FAME and ASE entries was counted to summarize the number of entries per scale and theme.

## Data analysis

To allow the same treatment of both quantitative and qualitative data, verbal counts from qualitative studies were translated into numerical results in the matrix. For example, adjectives deriving from the qualitative data such as 'never', 'sometimes', or 'always' or 'little', 'moderate', 'most', were transformed into categories with the numerical values '0', '0.5', or '1' [46–48]. Quantitative data were coded into the same numerical values according to the *P*-value and frequencies: *P* > .10 and 0–33.3% (value 0); *P* > .05 into *P*.10 and [Q5]33.4–66.6% (value 0.5); *P*.001 into *P*.05 and 67.7–100% (value 1). Numerical scoring entries were made when the feasibility, appropriateness, meaningfulness, and effectiveness of the ASE theme promoted the utility of MLC (value 1), had no effect on utility (value 0.5), or did not promote the utility of MLC (value 0). If a report did not address a certain theme, the cell was left blank. Assigning the level of utility was a process among four researchers (YK, JD, VB, ET).

For analysis, the matrix was exported into SPSS® version 28. To analyse a complete data set, multiple imputation was used for the missing values with the Markov Chain Monte Carlo (MCMC) method [49]. Two-sided Bayesian Pearson correlations were calculated to establish the strength of the relationship(s) between the FAME categories and the ASE themes and between the ASE themes[50,51]. It was hypothesized that the ASE themes correlate with the FAME themes (H). A Bayes Factor (BF) of >100 was considered as extreme evidence for H<sub>1</sub>, 30–100 as very strong evidence, 10–30 as strong evidence, 3–10 as moderate evidence, 1–3 as anecdotal evidence, 1 as no evidence, and a BF <1 was considered in favour of H<sub>0</sub> [51]. The criteria were set at 10,000 MC samples, a tolerance of 0.0001, and a maximum of 2000 iterations. This study is

considered as a first attempt in MLC behaviour framework building, lacking full-domain specific numerical information to facilitate good predictions and therefore we did not set priors for the analysis [31,52].

## Results

### Sample characteristics

The included 26 articles were published between 2010 and 2021. Most of the studies ( $n = 17$ ) had a qualitative design. The five quantitative studies all had a cross-sectional design and four studies a mixed method approach. Self-reported data from 4,785 midwives (MLC midwives  $n = 1,676$ /non-MLC midwives  $n = 3,109$ ) were retrieved. All but one study was conducted in the Oceanic countries ( $n = 16$ ) and in Europe ( $n = 9$ ) (in total 12 different countries). Midwives in the samples were either employed or self-employed and worked in different maternity care settings (e.g. hospital, community). The studies reported on various MLC strategies (e.g. caseload, team midwifery). Caseload midwifery was most often reported ( $n = 10$ ). The midwives in the studies cared for low-risk women ( $n = 10$ ), high-risk women ( $n = 1$ ), or for mixed groups of high and low-risk women ( $n = 14$ ). From one study, the risk status of women was unclear. Details of the studies are presented in Table 1.

**Table 1.** Papers included in the review.

Author(s), year of publication	Design study	Country	Aim of study	Population/midwives in the sample	FAME/ASE entries ( $n$ )
Clemons et al. [53]	Qualitative survey with open-ended questions	New Zealand	To explore job autonomy amongst midwives and to identify what supports/enables and what hinders/disables midwives' ability to practice autonomously within their role	253 midwives: 117 self-employed midwives, 88 employed and 48 'other'; 0–42 years of work experience; caring for low and high-risk women	2
Kashani et al. [54]	Qualitative, descriptive (interviews)	Australia, North-East Victoria	To describe midwives' experiences of and views on working in a CL midwifery model in a rural setting	11 qualified midwives; 24–61 years of age; 3–39 years of work experience; 5 presently and 6 midwives formerly practising in a CL model with low-risk women	18
Newton et al. [55]	Quantitative, cross-sectional (survey)	Australia	To compare views of CL midwifery: those working in CL models and those in SC models in hospitals with and without CL	542 midwives, working in public hospitals providing care to low-risk women: 20% CL midwives, 39% non-CL midwives, 41% were working in a hospital without a CoC model; 56% >10 years of work experience, 13% in the first year of practice	17
McInnes et al. [56]	Realist evaluation, participatory research, survey including open text answers, interviews, observations, audits, field notes	Scotland, UK	To explore how CoMC works, for whom, in what context, and to what extent, to inform sustainable on-going implementation and up scaling the CoMC model	Midwives working in one Scottish Health Board: baseline survey (321 midwives); interviews (113/77 CoMC/36 non-CoMC midwives); caring for low and high-risk women	14
Hollins Martin et al. [57]	Realist evaluation, online survey with free text comments	Scotland, UK	To collect baseline data prior to implementing a CoMC model collecting data about midwives' personal and professional wellbeing prior to service reorganization, with a longitudinal study intended to measure a change in midwives' reports across time <i>This paper reports the baseline data collection</i>	321 midwives working in a Scottish Health Board, including urban and rural settings with approx. 9500 births per year. The Health Board comprised of three settings: two hospitals and community midwifery services caring for low and high-risk women	12
Styles et al. [58]	Qualitative, implementation research (interviews)	Australia, Queensland	To explore the perceptions and experiences of midwifery and obstetric staff during the implementation and upscaling of midwifery CoC. Data collection within the first 2–3 months and 2 years after CoC implementation	15 midwives, core midwives, and midwives providing CoC to low and high-risk women <i>obstetric staff experiences excluded from the analysis</i>	11
Bradfield et al. [59]	Qualitative, descriptive phenomenological design, purposive sample (interviews)	West Australia	To explore the experiences of being 'with woman' during labour and birth from the perspectives of midwives working in a model where care is provided by a known midwife	10 midwives working in a 'known' midwife scheme; 35–57 years of age; 4–34 years of work experience; low and high-risk women. All midwives previously worked in a public hospital with fragmented care where the woman was unknown to them	12
Bradfield et al. [60]	Qualitative, descriptive phenomenological design (interviews)	West Australia	To explore midwives' perceptions of being with women during labour and birth in the context of various models of maternity care	31 midwives: 10 CoC midwives, 11 worked in a private obstetric-led model, 10 in standard public model; providing care to low and high-risk women; 35–62 years of age; 3–35 years of midwifery experience	8
Cronie et al. [61]	Quantitative, cross-sectional, online questionnaire	Netherlands	To measure and compare job satisfaction among hospital and primary care midwives	508 midwives, 103 hospital, and 405 primary care midwives; mean age 40 years; providing care to low-risk women	8

Taylor et al. [62]	Quantitative, cross-sectional (33 quantitative/16 qualitative questions)	United Kingdom	To examine the working patterns that midwives are willing and able to adopt, and to ascertain what barriers exist and what would help midwives to work in CoC models	798 midwives in 7 geographical areas (27 hospitals); providing care to low and high-risk women in community (36%) and obstetric units (34%); 57% had been qualified > 10 years	7
Dawson et al. [63]	Quantitative, cross-sectional (survey)	Australia	Comparing the experiences of CL and non-CL midwives in relation to burnout and attitudes to their professional role	542 midwives: 107 working in CL, 212 in other areas of midwifery in a hospital with a CL model, 220 worked in a hospital without a CL model; 56% had been practising as a midwife > 10 years. Risk status of women was unclear	5
Hunter et al. [64]	Qualitative, hermeneutic phenomenology (interviews)	New Zealand	To reveal what enables, safeguards, and sustains midwives to provide intrapartum care in freestanding midwifery-led units from midwives' and obstetricians' perspectives	4 CL community midwives, 7 employed midwives caring for low-risk women <i>non-midwives' responses excluded from analysis</i>	1
Coddington et al. [65]	Qualitative, descriptive exploratory study (interviews)	Australia	To examine midwives' experiences of transitioning from providing hospital-based midwifery to homebirth midwifery care	9 midwives and 4 midwifery managers (also practising) providing care to low-risk women in an MGP or community within the 5 past years	16
Dixon et al. [66]	Quantitative, cross-sectional (survey)	New Zealand	To describe and compare the demographic and work-related factors of midwives and to explore factors associated with burnout in groups of employed and self-employed midwives	473 self-employed midwives, 452 employed midwives, 148 both self-employed and employed midwives caring for low and high-risk women	7
Jepsen et al. [67]	Qualitative, phenomenology, (field observations, interviews)	North Denmark	To explore what constitutes CL midwifery from the perspectives of the midwives and how midwives experience working in CL midwifery	13 CL midwives; 2 different hospitals; 29–59 years of age; 4–25 years of work experience; 23% CL; 77% combination of CL and SC; caring for a mixed group of low and high-risk women	15
Newton et al. [68]	Mixed method (surveys and in-depth interviews)	Australia, Melbourne	To explore CL and SC midwives' views and experiences of midwifery work in two new CL models (survey at baseline and two years later; interviews at 6-months and after 2 years or at resignation)	Full- and part-time working CL (survey and interviews) and SC midwives (survey); 50% > 10 years of work experience; 21 CL and 130 SC midwives at baseline; 22 CL and 133 SC midwives at 2-years; 28 interviews with CL midwives; providing care to low-risk women	35
Bureau and Overgaard [69]	Qualitative multi-case design (interviews and focus groups)	Denmark	To explore: (1) The interplay between midwives and management in their negotiations on the introduction and development of CL midwifery (2) The professional and organizational interests pursued by midwives and management in the process (3) The professional and organizational resources activated by midwives and management in the process	(Deputy) Chief-, CL-, and ward midwives working in three hospitals (university, mid-level, community; low and high-risk women) in which CL midwifery had been introduced less than a year ago: 10 individual interviews (3 chief midwives, 3 deputy chief midwives, 3 CL midwives, 1 obstetrician); 14 group interviews (14 CL midwives, 10 ward midwives, 9 health visitors, 6 obstetricians) <i>non-midwives' responses excluded from analysis</i>	13
Cummins et al. [70]	Qualitative, descriptive (interviews)	Australia	To explore the experiences of the new graduate midwives who have worked in CoMC, to examine the support they received and, to establish the facilitators and barriers to the expansion of new graduate positions in midwifery CoMC	13 newly graduate midwives; 21–46 years of age; working within their first or second year of practice; caring for low and high-risk women	11
Maillefer et al. [71]	Qualitative, descriptive (interviews and focus groups)	Switzerland (Canton of Vaud)	To explore the perceptions of women and healthcare providers related to the future development of an MLC at a university hospital	10 midwives working at a university (tertiary) hospital caring for low and high-risk women <i>non-midwives' responses excluded from analysis</i>	4
Sjöblom et al. [72]	Qualitative, phenomenology (Interview)	Denmark, Iceland, Norway, Finland, Sweden	To describe the lived experience of being a homebirth midwife in the Nordic countries	21 homebirth midwives: 8 from Sweden, 5 from Denmark, 4 from Norway, 2 from Finland, and 2 from Iceland; 2–38 years of work experience; providing care to low-risk women	8

Warmelink et al. [73]	Qualitative, constructivist/interpretative design	Netherlands	To gain an understanding of how primary care midwives feel about their work and investigate factors associated with job satisfaction of primary care midwives	99 midwives; 22–61 years of age; 40 years of work experience; providing care to low-risk women	13
Edmondson and Walker [74]	Qualitative, grounded theory (interviews)	Australia, North Queensland	To uncover how birth centre midwives, working within midwifery CL, constructed their midwifery role to maintain a positive work-life balance	7 midwives; 0–3 years of work experience; 40–60 years of age without having young children; working at the birth centre with low-risk women	10
Menke et al. [75]	Qualitative, descriptive (focus groups)	Australia, Queensland	To examine midwives' perceptions of the organizational structures and processes of care and their impact on a CL model for socially disadvantaged women and vulnerable childbearing women	17 midwives working in MGP; 26–61 years of age; on average 8 years of work experience and 21 months of MGP experience; providing care to high-risk women	16
Newton et al. [20]	Quantitative, cross-sectional (survey)	Australia, Victoria	To compare midwives' attitudes to their professional role and measures of burnout between CL midwives and those working in SC models in these two newly introduced CL models	22 CL midwives and 133 SC midwives; CL midwives providing care to low-risk women; SC midwives providing care to a mixed group of low and high-risk women	4
Gu et al. [76]	Qualitative, phenomenology (interviews)	China, Shanghai	To explore and describe midwives' experiences of providing continuity of care to labouring women in a hospital setting	12 midwives; 3–8 years of work experience; providing care to a mixed group of low and high-risk women	10
Fereday and Oster [77]	Qualitative, interpretive study (Interviews)	Australia, Adelaide	To gain insight into the management of flexible work hours to achieve a work-life balance for a group of midwives working in a CL model	17 midwives, 3 months-2 years' experience in an MGP providing care to low-risk women	15

Continuity of Care (CoC)/Continuity of Midwifery Care (CoMC): a nominated primary care midwife who migrates with the woman in case of complications and/or referral of care.

Caseload (CL): one midwife providing all care, including being on-call for own caseload births with the backup of another midwife.

Midwifery Group Practice (MGP): a team of midwives sharing a caseload of women with backup from one or more midwives in the group, providing antenatal, intrapartum, and postnatal care in a scheduled system.

Standard Care (SC): care as practised in the place where the study was conducted (other than CoC/CoMC, CL, or MGP) such as shared care (hospital)midwife/obstetrician, (hospital)midwife/General Practitioner, care from a variety of midwives and obstetric staff, hospital care, care delivered by midwives who work shifts of 8–10 h.

## Quality

The JBI-criteria 'locating the researcher culturally or theoretically' ( $n = 11$ ) and 'is the influence of the researcher on the research, and vice-versa, addressed' ( $n = 10$ ) were recurrently unclear or absent in the qualitative methodological designs. Only one quantitative paper 'identified or stated confounding factors'. The papers showed overall good quality.

## FAME & ASE entries

In total, 292 entries were extracted from the data, with a minimum of one entry to a maximum of 35 entries per study (Table 1). 'Appropriateness' was the most often reported FAME theme and 'supportive factors' was the most often reported ASE theme (Table 2).

**Table 2.** Data matrix of entries per study.

Fame categories	Study	ASE themes											
		Attitude	Social influence (public)	Social influence (maternity care domain)	Self-efficacy	Regulation	Supportive factors	Barriers	Social norm	Professional norm	Personal norm	Knowledge	Intention
	Clemons et al. [53]			1				1					
	Newton et al. [55]					1	2	3					
	McInnes et al. [58]					1	2	2					
	Hollins Martin et al. [57]				1		1						
	Styles et al. [58]						1	1		1			

Feasibility N = 56	Bradfield et al. [59]					1				1			
	Coddington et al. [65]	1	1	1									
	Burau and Overgaard [69]	2				1	2	1		1			1
	Maillefer et al. [71]	1								1			
	Sjöblom et al. [72]	1	1							1			
	Warmelink et al. [73]					1	3	1	1	1	1	1	
	Edmondson and Walker [74]					1	3			1	1	1	
	Menke et al. [75]									1	1		
	Newton et al. [20]					1							
	Gu et al. [76]									1			
	Fereday and Oster [77]										1		
	N entries per study	<b>5</b>	<b>2</b>	<b>2</b>	<b>1</b>	<b>7</b>	<b>14</b>	<b>9</b>	<b>1</b>	<b>9</b>	<b>4</b>	<b>2</b>	<b>1</b>
Appropriateness N = 184	Kashani et al. [54]	3					8	1	1	2			1
	Newton et al. [55]	4	1	2			8	2	1	4	3	1	
	McInnes et al. [56]			1	1								
	Hollins Martin et al. [57]	4	1							1	2	1	
	Styles et al. [58]						1		2	2	2		
	Bradfield et al. [59]	1					2			1			
	Bradfield et al. [60]			1			4			3			1
	Cronie et al. [61]	1		1		3	3						
	Taylor et al. [62]						3	2		1		1	
	Hunter et al. [64]				1								
	Coddington et al. [65]	2	2			1	4			1	1	2	
	Dixon et al. [66]	1		1		3	1		1				
Jespen et al. [67]	2					2			3	1		1	

	Burau and Overgaard [69]						1		2				
	Cummins et al. [70]	2					6	1		1			
	Maillefer et al. [71]	1											
	Sjöblom et al. [72]						1	1					
	Warmelink et al. [73]								1				
	Edmondson and Walker [74]						1			1			
	Menke et al. [75]		1					4	4	1	1		2
	Newton et al. [20]	3				1	2	3			7		
	Gu et al. [76]						1	2		1	1		
	Fereday and Oster [77]						5	5		3			
	<i>N</i> entries per study	<b>24</b>	<b>5</b>	<b>6</b>	<b>2</b>	<b>8</b>	<b>53</b>	<b>21</b>	<b>12</b>	<b>25</b>	<b>18</b>	<b>5</b>	<b>5</b>
Meaningfulness <i>N</i> = 18	Kashani et al. [54]	1											
	Newton et al. [55]									1			
	McInnes et al. [56]											1	
	Hollins Martin et al. [57]									1			
	Styles et al. [58]												1
	Bradfield et al. [59]		1		1					1			
	Jespen et al. [67]	1							1				
	Maillefer et al. [71]	1											
	Sjöblom et al. [72]	1					1						1
	Warmelink et al. [73]									1			
	Edmondson and Walker [74]							1					
	Menke et al. [75]						1						
	Fereday and Oster [77]						1						
<i>N</i> entries per study	<b>4</b>	<b>1</b>	<b>-</b>	<b>1</b>	<b>-</b>	<b>3</b>	<b>1</b>	<b>1</b>	<b>4</b>	<b>-</b>	<b>1</b>	<b>2</b>	



Effectiveness N = 33	Kashani et al. [54]									1			
	Newton et al. [55]								3	1			
	McInnes et al. [56]							2		1	1	2	
	Bradfield et al. [59]						1		1				
	Dawson et al. [63]	1							1	1			1
	Jespen et al. [67]	1		1				1					1
	Newton et al. [68]	1								1			1
	Burau and Overgaard [69]	1											1
	Cummins et al. [70]						1						
	Warmelink et al. [73]	1								1			
	Gu et al. [76]				1			2				1	
N entries per study	5	–	1	1	–	1	4	2	5	6	2	6	
Total N entries per ASE theme	38	8	9	5	15	71	35	16	43	28	10	14	

### FAME themes

*Feasibility* of MLC was determined by childbearing women [60,65,72] and by leadership, staff, hours, equipment, rota, quality assurance in the midwife's work environment, and the extent this environment advocated MLC values [56,57,60,65,68–70,72,73]. *Feasibility* was shaped by midwives with MLC expertise in the culture of maternity services [20,53–56,58,65,68,69]. MLC was *feasible* when midwives were seen as knowledgeable and autonomous practitioners whilst collaborating on multi-disciplinary level [53,59,69,71–74,76]. MLC was *feasible* on an individual midwife level when a supportive network at home was available [57,61,62,68,70,73] and when midwives were intrinsically motivated, interested, and committed to provide MLC [69,71,72,76,78].

The woman-midwife relationship/partnership, working in a homely environment, continuity of care and carer, advocacy, autonomy, and empowerment were strategies of clinical *appropriateness* of MLC [57,59,60,65,67,71]. MLC was *appropriate* as it fitted the midwife's autonomous professional identity, centred around the strong midwife-woman relationship while providing continuous and individualized care [54,57,61,67,68]. MLC was *appropriate* in the context of informed and shared decision-making in maternity services [65,66,78].

MLC was determined to be *meaningful* when midwives believed in women and when women believed in themselves – in the ability to give birth [54,72,75]. *Meaningfulness* was shown by women trusting midwives and midwives being true to midwifery values and beliefs [54,56,57,68,72,74]. It was *meaningful* for midwives to know the women in their care, to engage with them and their families, and with women's individual circumstances and needs [56,67,73]. Continuity was *meaningful* in the relationship between the midwife and the woman [54,57,69–71]. Midwives agreed that the benefits of MLC were worth the efforts and therefore *meaningful* [58].

The *effectiveness* of MLC was recognized by increased satisfaction of women with care [56], job satisfaction of midwives, and lower levels of burnout compared to non-MLC midwives [56,63,69]. Midwives reported MLC *aseffective* because of observed and experienced improved medical and psychosocial standards of care and improved clinical outcomes for birthing women and their babies [56,68,69].

### ASE themes

MLC midwives showed a positive *attitude* towards the model of care [56,63,67,68] and regarded MLC as important for (skills) development, expertise, and professional autonomy [54,57,65,66,68,69,71,73] and to be 'a good midwife' [54,56,68]. Midwives believed that MLC could be the opportunity to introduce change in the medical model of maternity services [71]. The midwives had a fundamental belief that MLC improved the woman's ability to give birth and to improve birth outcomes [68,72].

*Social influence* was observed within maternity services (colleagues) as well as in the public domain (women, media) [54,56,61,65,66,68]. Midwives reported that their role of being the woman's advocate was often met with resistance, disrespect, and scrutiny or questions from obstetricians, midwives [53,56,60,65,68], and from colleagues – who believed MLC exposes mother and baby to potential health risks [72]. Midwives were affected by negative MLC media messages and hospital management's announcements [65], feeling unsupported and judged [53,75].

*Self-efficacy* was a result of strong individual MLC values and beliefs, reliance on skills [64], and feeling empowered and confident [53,56,65,68]. Midwives indicated that through developing a close and trusting relationship with the woman, knowing the woman's health and her family situation, they believed in their abilities to provide optimal care for women [54,70,75,76].

Not being able to work within a woman-centred care approach was perceived as a *barrier* to construct the MLC role [56]. Poor workforce engagement of colleagues, poor leadership, lack of regulation, poor management, and/or miscommunications with the wider healthcare team [53,56,65,66,69,73,75], a lack of staff [70] and activities such as meetings and paperwork were *barriers* in adopting MLC [54,73]. Unclear/undefined working hours were *barriers* to achieve work-life balance [20,57,58,68,72,76,77]. For non-MLC midwives, being on-call was the primary reason not to work in the MLC model [20,68]. Additionally, some midwives regarded having young children and having to organize childcare as *barriers* for being an MLC midwife [62,68,77].



Feasibility (F)	1.11	0.00	0.75	<b>37.92</b>	20.1	0.05	1.50	0.00	16.58	9.68	26.20	<b>38.15</b>
Appropriateness (A)	<b>41.29</b>	<b>32.06</b>	0.09	27.44	0.00	20.48	<b>37.40</b>	0.93	<b>37.67</b>	5.22	2.72	11.22
Meaningfulness (M)	1.32	6.40	0.00	2.87	0.00	<b>31.10</b>	2.91	0.00	20.65	0.05	8.44	0.00
Effectiveness (E)	23.66	20.41	0.00	17.56	<b>41.48</b>	11.95	1.40	3.07	4.71	<b>38.13</b>	<b>39.59</b>	0.00
Attitude		22.30	0.58	<b>33.79</b>	0.00	11.20	<b>36.24</b>	<b>38.16</b>	2.17	0.76	<b>39.87</b>	0.86
Social Influence public			<b>41.48</b>	<b>41.18</b>	<b>40.88</b>	27.84	0.03	24.09	<b>38.93</b>	0.92	19.85	0.08
Social Influence maternity domain				<b>41.55</b>	0.00	<b>32.82</b>	19.60	<b>38.08</b>	5.61	1.63	0.00	0.00
Self-efficacy					1.19	8.43	5.24	0.72	15.05	2.08	4.37	0.50
Regulation						5.67	0.45	0.05	0.96	<b>41.59</b>	0.00	<b>41.58</b>
Barriers							28.52	19.51	17.95	3.36	<b>40.42</b>	<b>40.41</b>
Supportive factors								27.24	12.16	0.00	25.62	<b>40.85</b>
Social norm									0.02	20.15	0.00	1.12
Professional norm										26.57	2.64	<b>31.86</b>
Personal norm											3.78	22.97
Intention												1.55

Bayes Factor: BF >100 = extreme evidence for H<sub>1</sub>; **BF 30–100 = very strong evidence H<sub>1</sub>**; BF 10–30 = strong evidence for H<sub>1</sub>; BF 3–10 = moderate evidence for H<sub>1</sub>; BF 1–3 = anecdotal evidence for H<sub>1</sub>; BF 1 = no evidence for H<sub>1</sub>; BF <1 = extreme evidence for H<sub>0</sub>.

## Discussion

The framework of the FAME scales and the themes of the extended ASE model showed that the utility of MLC is explained and influenced by multiple and diverse midwives' behavioural factors. The various behavioural determinants of midwives related to the utility of MLC were entangled in a systematic way, which has not been done before. The predominantly strong and very strong levels of evidence and to a lesser degree the moderate and anecdotal evidence support the hypothesis that ASE themes correlate with the FAME scales.

The data entries most often reported on MLC in the midwifery context and culture (appropriateness scale), the results strongly emphasize that MLC seems to be regarded as a core and true value of midwifery [23]. However, the thematic findings show that the MLC context and culture can be distinguished in prosocial intrinsic (attitude, professional norm) and extrinsic pragmatic forces (supportive factors, social influence of the public) [81] – suggesting an ambiguity in forces [82]: Some midwives value MLC because it fosters own happiness and wellbeing as well as of the women in their care. MLC also aligns with the core value of 'doing or being good' [83]. Other midwives value MLC because of having sufficient time to provide care, a reduced caseload, a fixed team structure, and autonomous and flexible diary management [84]. Both prosocial and pragmatic factors are important for MLC midwives' self-determination and their resilience [81] as these factors contribute to midwives' competency, autonomy, and relatedness to working in the MLC model [82]. Based on the prosocial-pragmatic duality, midwives should be supported to be unique in providing MLC and not utilize a one-size-fits-all MLC approach, accepting and recognizing that midwives have different perceptions to provide MLC and interact differently with the women in their care [19,41,79]. There should be no preference for either the prosocial or the pragmatic aspects – although usually the prosocial aspects precede the pragmatic aspects and then co-exist [85]. Additionally, as different MLC strategies exist, it might be that some strategies (e.g. group practice) suit some midwives better in achieving a family–work balance [41,62,68,77]. In the woman–midwife relationship, the midwife is as much an individual as the woman [85]. To avoid interdisciplinary polarization or dis cohesion, reasons attributing to midwives' self-determination and their resilience need to be considered, acknowledged, and respected, as well as it needs, therefore, to be accepted that MLC might not fit all midwives [24,40,41,84,85].

The findings enhance the predominant monodisciplinary midwifery perspective embedded in the midwifery culture. Midwives did not often report on MLC in the wider context of integral maternity care (feasibility) while MLC exists within a broad cultural, socio-political, and financial climate and multi-disciplinary network of health and maternity services [1,86,87]. The analysis shows that feasibility is predominantly underpinned by self-efficacy, professional norm, intention, and knowledge. The thematic findings about these behavioural aspects do not include referral to the multi-disciplinary maternity service culture – instead, these are reports on a midwife–woman level. Midwives' limited reports on MLC feasibility within the maternity services climate might be because of the feeling of the patriarchal and hierarchical medical hegemonic and medical dominant maternity care system with multi-disciplinary debates about professional responsibilities and abilities [9,24,80,86–88]. Nevertheless, for increased feasibility of MLC, while reclaiming the midwifery role in a medicalized climate, midwives will need to collaborate effectively in the organizational climate of maternity services and to stand up for their professional identity in hierarchal relationships [89]. However, midwives are not able to achieve increased feasibility as an individual or separate monodisciplinary group. Therefore, while focusing on multi-disciplinary collaboration, simultaneously greater insight, reflection on traditional power imbalances, and respect for each other's job (autonomy) should be established to adequately embed MLC in maternity services [90].

Evaluation of MLC consistently demonstrates quality of care, good outcomes for mothers and babies, and overall satisfaction of maternity experiences for women. For MLC to be (come) regarded as a first choice for all childbearing women, it seems imperative that the worth and effectiveness of MLC should be shared, internalized, and formalized in the maternity climate to avoid the domination of obstetricians' regulation in the maternity care climate [90]. It is evident that practice transformation needs to be supported through theoretical understanding grounded in contextual significance to implement MLC as a sustainable model of care and handing a mandate to MLC midwives to assist organizational change and to prevent medical/obstetric care to have a head start in the feasibility of MLC [91]. MLC needs an emancipatory shift and prominent change agents and role models, transformational leaders who provide vision and resources, and an active coordination of knowledge exchange on meso-level [87]. MLC needs academics, policymakers, and governors in midwifery organizations, putting MLC up in a debate with healthcare politicians and health services/insurances on macro level [24]. Effective and efficient educational programs to improve (student)midwives' knowledge during shifting the context of (re)orientation to MLC and to grow confidence would be of merit [85,92]. Additionally, midwives need to find a way to (re)claim their legal right to act autonomously and reflect on their midwifery abilities by using strategies primarily consisting of self-awareness and self-determination [59,76]. The qualitative findings show that the woman–midwife relationship is an important and recurring issue. Therefore, it might be necessary to seek an alliance with women as MLC key stakeholders towards pushing the change at the social level to indicate for MLC to

become politically and financially governed [83].

The analysis shows extreme evidence that building meaningful trusting relationships with women and the commitment to the physiological approach of childbirth are not correlated with the influence of the maternity care domain, regulation, the midwife's social and personal norms, and/or with the midwife's knowledge [54,56,57,60,67,68,72–75]. The thematic findings suggest that MLC midwives are not so much being put off by human and non-human barriers in the work setting or by issues in their personal lives. The meaningfulness of MLC and midwives' MLC actions are seemingly separate entities, emphasizing the epistemological status of MLC [41,85,93]. It therefore must be safeguarded for the barriers in the work setting to become conditions that affect the meaningfulness of MLC [94]. Constant and conscious reflection on the meaningfulness of MLC among (student)midwives is of merit.

## Limitations

Although several behavioural determinants were identified, this does not assure the strength of the ASE model in explaining the utility of MLC. There was an inconsistency in the number of reported FAME and ASE attributes, causing over and underrepresentation of certain themes. Over and underrepresentation could have been a result of including studies performed in the same setting and same group of researchers, reporting on the same issues, likely to contribute to publication bias. Having included reports of both MLC and non-MLC midwives limits potential positivity and/or negativity reporting bias [84]. The imputation of missing data could also have added to publication bias, however, using the MCMC algorithm reduced the loss of precision [95]. The lack of evidence/limited number of entries in certain areas (e.g. meaningfulness, self-efficacy, social norm, intention) can be considered as topics for future research to complement the framework that was developed based on the current evidence. From all the Bayes Factors, there were no BFs showing extreme evidence, and the BFs showing very good evidence had a ceiling of 41.6. Multiple determinants showed no evidence to test the hypothesis. However, using a Bayesian approach for estimation of  $H_1$  provides outcomes in a form that is less prone to misinterpretation, aiding to the trustworthiness of the final framework [51,96]. The included studies showed heterogeneity of sample setting characteristics. Most of the included studies were conducted in Australia and Europe with midwives – potentially overrepresenting and underrepresenting certain geographical areas, midwifery cultures, maternity settings, and health care systems, affecting the generalizability of the results. As MLC was first introduced in predominantly New Zealand, Australia, and the United Kingdom, it might not be surprising that most studies originated from these countries [85]. Not all studies provided information about midwives' backgrounds such as years of experience, likely to cause confounding. Additionally, a variety of different strategies of MLC were reported. The study did not address the variations between the studied geographic distributions, midwives' characteristics, or the different organizational settings or care strategies, possibly limiting the interpretation of the findings. Sub analyses can be recommended for future research. The fact that there was sometimes weak to no evidence of relations between FAME and ASE themes gives rise to the thought that other variables and/or unknown confounders not studied yet may affect midwives' behaviour. Hence, further research is recommended. Although the studies showed overall good quality, there were unclear or absent JBI-criteria, lowering the overall appraisal. JBI has no clear guidelines about exclusion [39,40] and the decision was based on discussion among the researchers, which could have led to selection bias based on the authors' beliefs about MLC [97].

## Conclusion

The body of evidence regarding safety and efficacy of MLC is well established. The study provides a theoretical underpinning, offering a grass roots approach to practice grounded in contextual significance. The findings and framework are a first attempt to structure the midwives' behavioural determinants related to the utility of MLC, offering recommendations for the implementation, scale up, and sustainability of this model of care. MLC is without a doubt a core and true value of midwifery, albeit not being fully embraced by all midwives. To implement and maintain MLC a multipronged approach is needed, with attention for the strong and very strong behavioural aspects of midwives related to the utility of MLC. It needs to be considered how these aspects are addressed in practice, multi-disciplinary collaboration, education, management, and/or policies and guidelines of antenatal, intrapartum, and postnatal care. The evidence indicates the need for an emancipatory transition for MLC to become fully embedded in the wider culture and climate of maternity services for MLC to be (come) the first choice for all childbearing women. This study offers opportunities for future research to update and/or complement the framework that was developed.

## Disclosure statement

No potential conflict of interest was reported by the author(s) [Q7].

## ORCID

Yvonne Kuipers <http://orcid.org/0000-0002-4200-0522>

Julie Degraeve <http://orcid.org/0000-0002-9971-4622>

Valerie Bosmans <http://orcid.org/0000-0002-6802-4485>

Ellen Thael <http://orcid.org/0000-0002-0651-6652>

Eveline Mestdagh <http://orcid.org/0000-0003-0116-0738>

## References [Q8]

- 1 Sandall J, Soltani H, Gates S, et al. Midwife-led continuity models versus other models of care for childbearing women. *Cochrane Database Syst Rev* C. 2016;4(4):d004667.
- 2 Dixon L, Prileszky G, Guillard K, et al. What evidence supports the use of free-standing midwifery led units (primary units) in New Zealand. *N Z Coll Midwives*. 2012;46:13–20.
- 3 McLachlan H, Forster D, Davey M-A. A randomised controlled trial of caseload midwifery for women at low risk of medical complications (COSMOS): women's satisfaction with care. *J Paediatr Child Health*. 2012;48:8–81.
- 4 Allen J, Gibbons K, Beckmann M, et al. Does model of maternity care make a difference to birth outcomes for young women? A retrospective cohort study. *Int J Nurs Stud*. 2015;52(8):1332–1342.
- 5 McLachlan HL, Forster DA, Davey M-A, et al. The effect of primary midwife-led care on women's experience of childbirth: results from the COSMOS randomised controlled trial. *Bjog*. 2016;123(3):465–474.
- 6 Miller S, Abalos E, Chamillard M, et al. Beyond too little, too late and too much, too soon: a pathway towards evidence-based, respectful maternity care worldwide. *Lancet*. 2016;388(10056):2176–2192.
- 7 Shaw D, Guise J-M, Shah N, et al. Drivers of maternity care in high-income countries: can health systems support woman-centred care? *Lancet*. 2016;388(10057):2282–2295.
- 8 Homer C, Leap N, Edwards N, et al. Corrigendum to "Midwifery continuity of carer in an area of high socio-economic disadvantage in London: a retrospective analysis of Albany Midwifery Practice outcomes using routine data (1997–2009)" [Midwifery 48 (2017) 1–10]. *Midwifery*. 2017;51:53.
- 9 McInnes RJ, Collins Martin CJ, MacArthur J. Midwifery continuity of carer: developing a realist evaluation framework to evaluate the implementation of strategic change in Scotland. *Midwifery*. 2018;66:103–110.
- 10 Gidaszewski B, Khajehei M, Gibbs E, et al. Comparison of the effect of caseload midwifery program and standard midwifery-led care on primiparous birth outcomes: a retrospective cohort matching study. *Midwifery*. 2019;69:10–16.

- 11 **Renfrew MJ, McFadden A, Bastos MH, et al.** Midwifery and quality care: findings from a new evidence-informed framework for maternal and newborn care. *Lancet*. 2014;384(9948):1129–1145.
- 12 **Rocca-Ihenacho L, Batinelli L, Thael S, et al.** Midwifery unit standards. London: London City University; 2018.
- 13 **Yu S, Fiebig DG, Scarf V, et al.** Birth models of care and intervention rates: the impact of birth centres. *Health Policy*. 2020;124(12):1395–1402.
- 14 **Macfarlane AJ, Blondel B, Mohangoo AD, et al.** Wide differences in mode of delivery within Europe: risk-stratified analyses of aggregated routine data from the Euro-Peristat study. *BJOG*. 2016;123(4):559–568.
- 15 **Homer C.** Models of maternity care: evidence for midwifery continuity of care. *MJA*. 2016;205(8):370–374.
- 16 **Symon A, Pringle J, Cheyne H, et al.** Midwifery-led antenatal care models: mapping a systematic review to an evidence-based quality framework to identify key components and characteristics of care. *BMC Pregnancy Childbirth*. 2016;16(168).[Q9]
- 17 **Kuipers Y, van Beeck E, van den Berg L, et al.** The comparison of the interpersonal action component of woman-centred care reported by healthy pregnant women in different sized practices in the Netherlands: a cross-sectional study. *Women Birth*. 2021;34(4):e376–e383.
- 18 **Homer C, Leap N, Brodie P, et al.** Midwifery continuity of care. 2nd ed. Chatswood: Elsevier; 2019.
- 19 **Michel-Schuldt M, McFadden A, Renfrew M, et al.** The provision of midwife-led care in low-and middle-income countries: an integrative review. *Midwifery*. 2020;84:102659.
- 20 **Newton M, Dawson K, Forster D, et al.** Midwives' views of caseload midwifery – comparing the caseload and non-caseload midwives' opinions. A cross-sectional survey of Australian midwives. *Women Birth*. 2021;34(1):e47–e56.
- 21 **Callander EJ, Slavin V, Gamble J, et al.** Cost-effectiveness of public caseload midwifery compared to standard care in an Australian setting: a pragmatic analysis to inform service delivery. *Int J Qual Health Care*. 2021;33(2).[Q10]
- 22 **Hewitt L, Dahlen H, Hartz D, et al.** Leadership and management in midwifery-led continuity of care models: a thematic and lexical analysis of a scoping review. *Midwifery*. 2021;98:102986.
- 23 **ICM.** Position Statement Midwifery led care, the first choice for all women (PS2011\_012 V2017 ENG). 2011. The Hague, Netherlands: International Confederation of Midwives (revised Toronto 2017).
- 24 **Sidebotham M, Fenwick J, Rath S, et al.** Midwives' perceptions of their role within the context of maternity service reform: an appreciative inquiry. *Women Birth*. 2015;28(2):112–120.
- 25 **Green LW, Kreuter MW.** Health program planning: an educational and ecological approach. 4th ed. New York (NY): McGraw-Hill; 2005.
- 26 **Eccles MP, Grimshaw JM, MacLennan G, et al.** Explaining clinical behaviors using multiple theoretical models. *Implement Sci*. 2012;7:99.
- 27 **Barkimer J.** Clinical growth: an evolutionary concept analysis. *ANS Adv Nurs Sci*. 2016;39(3):E28–E39.
- 28 **Bayes S, Fenwick J, Jennings D.** Readiness for practice change: evaluation of a tool for the Australian midwifery context. *Women Birth*. 2016;29(3):240–244.
- 29 **Salmon J, Hesketh KD, Arundell L, et al.** Changing behavior using ecological models. In: **Hagger M, Cameron L, Hamilton K, et al**, editors. *The handbook of behavior change*. Cambridge: Cambridge University Press; 2020:1-14.[Q11]
- 30 **Lopes Costa P, Graça AM, Marques-Quinteiro P, et al.** Multilevel research in the field of organizational behavior: an empirical look at 10 years of theory and research. *SAGE Open*. 2013;3(3):2158244013498244.
- 31 **Schoot R, Depaoli S.** Bayesian analyses: where to start and what to report. *European Health Psychologist*. 2014;16:75–84.
- 32 **Sandelowski M, Voils CI, Barroso J.** Defining and designing mixed research synthesis studies. *Res Sch*. 2006;13(1):29.
- 33 **Pearson A, Wiechula R, Court A, et al.** The JBI model of evidence-based healthcare. *Int J Evid Based Healthc*. 2005;3(8):207–215.
- 34 **ICM.** Contact details ICM Members. 2021. Available from: <https://www.internationalmidwives.org/about-us/membership/>
- 35 **Page MJ, McKenzie JE, Bossuyt PM, et al.** The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi:10.1136/bmj.n71
- 36 **Lockwood C, Munn Z, Porritt K.** Qualitative research synthesis: methodological guidance for systematic reviewers utilizing meta-aggregation. *Int J Evid Based Healthc*. 2015;13(3):179–187.
- 37 **Moola S, Munn Z, Tufanaru C, et al.** Chapter 7: systematic reviews of etiology and risk. In: **Aromataris E, Munn Z**, editors. *Joanna Briggs Institute reviewer's manual*. 2020. Available from <https://synthesismanual.jbi.global>. <https://doi.org/10.46658/JBIMES-20-08>
- 38 **Bartholomew Eldridge L, Markham C, Ruiter R, et al.** Planning health promotion programs: an intervention mapping approach. 4th ed. Hoboken (NJ): Wiley; 2016.
- 39 **Fontein-Kuipers YJ, Budé L, Ausems M, et al.** Dutch midwives' behavioural intentions of antenatal management of maternal distress and factors influencing these intentions: an exploratory survey. *Midwifery*. 2014;30(2):234–241.
- 40 **Kuipers Y, Boele A, Stuij C.** Midwives' perceptions of influences on their behaviour of woman-centered care: a qualitative study. *Front Wom Health*. 2016;1:20–26.
- 41 **Fontein-Kuipers Y, de Groot R, van Beeck E, et al.** Dutch midwives' views on and experiences with woman-centred care – a Q-methodology study. *Women Birth*. 2019;32(6):e567–ee75.
- 42 **Clarke V, Braun V, Hayfield N.** Thematic analysis. In: *Qualitative psychology. A practical guide to research methods*. 3rd ed. Los Angeles (CA): SAGE; 2015.[Q12]
- 43 **Armitage CJ, Conner M.** Efficacy of the theory of planned behaviour: a meta-analytic review. *Br J Soc Psychol*. 2001;40(Pt 4):471–499.
- 44 **Kwon HR, Silva EA.** Mapping the landscape of behavioral theories: systematic literature review. *J Plan Lit*. 2020;35(2):161–179.
- 45 **Corbin JM, Strauss AL.** Basics of qualitative research: techniques and procedures for developing grounded theory. 4th ed. Los Angeles (CA): SAGE; 2015.
- 46 **Voils C, Hasselblad V, Crandell J, et al.** A Bayesian method for the synthesis of evidence from qualitative and quantitative reports: the example of antiretroviral medication adherence. *J Health Serv Res Policy*. 2009;14(4):226–233.
- 47 **Crandell JL, Voils C, Chang Y, et al.** Bayesian data augmentation methods for the synthesis of qualitative and quantitative research findings. *Qual Quant*. 2011;45(3):653–669.
- 48 **Pearson A, Weeks S, Stern S.** Translation science and the JBI model of evidence-based healthcare. Philadelphia (PA): Lippincott Williams & Wilkins; 2011.
- 49 **Ma Z, Chen G.** Bayesian methods for dealing with missing data problems. *J Korean Stat Soc*. 2018;47(3):297–313.
- 50 **Matzke D, Ly A, Selker R, et al.** Bayesian inference for correlations in the presence of measurement error and estimation uncertainty. *Collabra Psychol*. 2017;3(1):25.
- 51 **Nuzzo RL.** An introduction to Bayesian data analysis for correlations. *PM&R*. 2017;9(12):1278–1282.
- 52 **Gelman A, Simpson DP, Betancourt M.** The prior can often only be understood in the context of the likelihood. *Entropy*. 2017;19:555.

- 53 **Clemons JH, Gilkison A, Mharpara TL, et al.** Midwifery job autonomy in New Zealand: I do it all the time. *Women Birth.* 2021;34(1):30–37.
- 54 **Kashani A, Ingberg JL, Hildingsson I.** Caseload midwifery in a rural Australian setting: a qualitative descriptive study. *Eur J Midwifery.* 2021;5:2.
- 55 **Newton MS, McLachlan H, Willis KF, et al.** Comparing satisfaction and burnout between caseload and standard care midwives: findings from two cross-sectional surveys conducted in Victoria, Australia. *BMC Pregnancy Childbirth.* 2014;14:426.
- 56 **McInnes RJ, Aitken-Arbuckle A, Lak S, et al.** Implementing continuity of midwife carer – just a friendly face? A realist evaluation. *BMC Health Serv Res.* 2020;20(1):304.
- 57 **Hollins Martin CJ, MacArthur J, Martin CR, et al.** Midwives' views of changing to a Continuity of Midwifery Care (CMC) model in Scotland: a baseline survey. *Women Birth.* 2020;33(5):e409–e419.
- 58 **Styles C, Kearney L, George K.** Implementation and upscaling of midwifery continuity of care: the experience of midwives and obstetricians. *Women Birth.* 2020;33(4):343–351.
- 59 **Bradfield Z, Hauck Y, Kelly N, et al.** Urgency to build a connection: midwives' experiences of being 'with woman' in a model where midwives are unknown. *Midwifery.* 2019;69:150–157.
- 60 **Bradfield Z, Hauck Y, Duggan R, et al.** Midwives' perceptions of being 'with woman': a phenomenological study. *BMC Pregnancy Childbirth.* 2019;19(1):363.
- 61 **Cronie D, Perdok H, Verhoeven C, et al.** Are midwives in the Netherlands satisfied with their jobs? A systematic examination of satisfaction levels among hospital and primary-care midwives in the Netherlands. *BMC Health Serv Res.* 2019;19(1):832.
- 62 **Taylor B, Cross-Sudworth F, Goodwin L, et al.** Midwives' perspectives of continuity based working in the UK: a cross-sectional survey. *Midwifery.* 2019;75:127–137.
- 63 **Dawson K, McLachlan H, Willis KF, et al.** Comparing caseload and non-caseload midwives' burnout levels and professional attitudes: a national, cross-sectional survey of Australian midwives working in the public maternity system. *Midwifery.* 2018;63:60–67.
- 64 **Hunter M, Smythe E, Spence D.** Confidence: fundamental to midwives providing labour care in freestanding midwifery-led units. *Midwifery.* 2018;66:176–181.
- 65 **Coddington R, Catling C, Homer CS.** From hospital to home: Australian midwives' experiences of transitioning into publicly-funded homebirth programs. *Women Birth.* 2017;30(1):70–76.
- 66 **Dixon L, Guillard K, Pallant J, et al.** The emotional wellbeing of New Zealand midwives: comparing responses for midwives in caseloading and shift work settings. *N Z Coll Midwives J.* 2017;53:5–14.
- 67 **Jepsen I, Mark E, Aagaard Nøhr E, et al.** A qualitative study of how caseload midwifery is constituted and experienced by Danish midwives. *Midwifery.* 2016;36:61–69.
- 68 **Newton MS, McLachlan H, Forster D, et al.** Understanding the 'work' of caseload midwives: a mixed-methods exploration of two caseload midwifery models in Victoria, Australia. *Women Birth.* 2016;29(3):223–233.
- 69 **Burau V, Overgaard C.** Caseload midwifery as organisational change: the interplay between professional and organisational projects in Denmark. *BMC Pregnancy Childbirth.* 2015;15:121.
- 70 **Cummins AM, Denney-Wilson E, Homer CS.** The experiences of new graduate midwives working in midwifery continuity of care models in Australia. *Midwifery.* 2015;31(4):438–444.
- 71 **Maillefer F, de Labrusse C, Cardia-Vonèche L, et al.** Women and healthcare providers' perceptions of a midwife-led unit in a Swiss university hospital: a qualitative study. *BMC Pregnancy Childbirth.* 2015;15:56.
- 72 **Sjöblom I, Lundgren I, Idvall E, et al.** Being a homebirth midwife in the Nordic countries – a phenomenological study. *Sex Reprod Healthc.* 2015;6(3):126–131.
- 73 **Warmelink JC, Hoijtink K, Noppers M, et al.** An explorative study of factors contributing to the job satisfaction of primary care midwives. *Midwifery.* 2015;31(4):482–488.
- 74 **Edmondson MC, Walker SB.** Working in caseload midwifery care: the experience of midwives working in a birth centre in North Queensland. *Women Birth.* 2014;27(1):31–36.
- 75 **Menke J, Fenwick J, Gamble J, et al.** Midwives' perceptions of organisational structures and processes influencing their ability to provide caseload care to socially disadvantaged and vulnerable women. *Midwifery.* 2014;30(10):1096–1103.
- 76 **Gu C, Zhang Z, Ding Y.** Chinese midwives' experience of providing continuity of care to labouring women. *Midwifery.* 2011;27(2):243–249.
- 77 **Fereday J, Oster C.** Managing a work-life balance: the experiences of midwives working in a group practice setting. *Midwifery.* 2010;26(3):311–318.
- 78 **Bradfield Z, Hauck Y, Kelly M, et al.** "It's what midwifery is all about": Western Australian midwives' experiences of being 'with woman' during labour and birth in the known midwife model. *BMC Pregnancy Childbirth.* 2019;19(1):29.
- 79 **Van den Branden L, Van de Craen N, Van Leugenhaege L, et al.** Flemish midwives' perspectives on supporting women during the transition to motherhood – a Q-methodology study. *Midwifery.* 2022;105:103213.
- 80 **Cronie D, Rijnders M, Jans S, et al.** How good is collaboration between maternity service providers in the Netherlands? *J Multidiscip Healthc.* 2019;12:21–30.
- 81 **Crowther S, Hunter B, McAra-Couper J, et al.** Sustainability and resilience in midwifery: a discussion paper. *Midwifery.* 2016;40:40–48.
- 82 **Freenny Y, Fellenz MR.** Work engagement as a key driver of quality of care: a study with midwives. *J Health Organ Manag.* 2013;27(3):330–349.
- 83 **Kuipers Y.** Exploring the uses of virtues of woman-centred care: a quest, synthesis and reflection. *Nurs Philos.* 2022; 23(2):e12380.[Q13]
- 84 **Dharni N, Essex H, Bryant MJ, et al.** The key components of a successful model of midwifery-led continuity of carer, without continuity at birth: findings from a qualitative implementation evaluation. *BMC Pregnancy Childbirth.* 2021;21(1):205.
- 85 **Fontein-Kuipers Y, de Groot R, van Staa A.** Woman-centered care 2.0: bringing the concept into focus. *Eur J Midwifery.* 2018;2:5.
- 86 **Perdok H, Jans S, Verhoeven C, et al.** Opinions of maternity care professionals and other stakeholders about integration of maternity care: a qualitative study in the Netherlands. *BMC Pregnancy Childbirth.* 2016;16(1):188.
- 87 **Pace C.** Co-creation of guiding principles and a practical framework for a midwifery continuity of carer model: a cooperative inquiry with midwives. Aberdeen: RGU; 2019.
- 88 **Jefford E, Jomeen J, Wallin M.** Midwifery abdication – is it acknowledged or discussed within the midwifery literature: an integrative review. *Eur J Midwifery.* 2018;2:6.
- 89 **Thompson SM.** Increasing self-efficacy in student midwives for physiological childbirth. Thesis. Maastricht University. 2020.
- 90 **van der Lee N, Driessen EW, Scheele F.** How the past influences interprofessional collaboration between obstetricians and midwives in the Netherlands: findings from a secondary analysis. *J Interprof Care.* 2016;30(1):71–76.
- 91 **Jacobs G, van Witteloostuijn A, Christe-Zeyse J.** A theoretical framework of organizational change. *J Organ Chang Manag.* 2013;26(5):772–792.
- 92 **Batinelli L, Thaelis E, Leister N, et al.** What are the strategies for implementing primary care models in maternity? A systematic review on midwifery units. *BMC Pregnancy Childbirth.* 2022;22.[Q14]
- 93 **Eri TS, Berg M, Dahl B, et al.** Models for midwifery care: a mapping review. *Eur J Midwifery.* 2020;4:30.

- 94 **Hultin L, Mähring M.** How practice makes sense in healthcare operations: studying sensemaking as performative, material-discursive practice. *Hum Relat.* 2017;70(5):566–593.
- 95 **Green PL, Worden K.** Bayesian and Markov chain Monte Carlo methods for identifying nonlinear systems in the presence of uncertainty. *Philos Trans A Math Phys Eng Sci.* 2015;373:2051.
- 96 **Van Doorn J, Ly A, Marsmn M, et al.** Bayesian Inference for Kendall's rank correlation coefficient. *Am Stat.* 2018;72:1537-2731.**[Q15]**
- 97 **Palaganas EC, Sanchez M, Molintas V, et al.** Reflexivity in qualitative research: a journey of learning. *Qualitative Report.* 2017;22(2):426–438.